

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrDOVATO

dolutegravir and lamivudine tablets

50 mg dolutegravir (as dolutegravir sodium) and 300 mg lamivudine, Oral

Antiretroviral Agent

ViiV Healthcare ULC
245, boulevard Armand-Frappier
Laval, Quebec
H7V 4A7

Date of Initial Approval:
August 22, 2019

Date of revision:
September 15, 2021

Submission Control No: 245212

©2021 ViiV Healthcare group of companies or its licensor
Trademarks are owned by or licensed to the ViiV Healthcare group of companies

RECENT MAJOR LABEL CHANGES

Contraindications (2)	02/2020
Warnings and Precautions, Pregnant Women (7.1.1)	06/2021
Warnings and Precautions, Renal (7)	08/2021
Dosage and Administration, Dosing Considerations (4.1)	10/2020
Dosage and Administration, Recommended Dose and Dosage Adjustment (4.2)	10/2020

TABLE OF CONTENTS

RECENT MAJOR LABEL CHANGES	2
TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	4
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations.....	5
4.2 Recommended Dose and Dosage Adjustment.....	5
4.3 Missed Dose	6
5 OVERDOSAGE	6
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	6
7 WARNINGS AND PRECAUTIONS	7
7.1 Special Populations.....	10
7.1.1 Pregnant Women.....	10
7.1.2 Breast-feeding	11
7.1.3 Pediatrics	12
7.1.4 Geriatrics	12
8 ADVERSE REACTIONS	12
8.1 Adverse Reaction Overview	12
8.2 Clinical Trial Adverse Reactions.....	12
8.3 Less Common Clinical Trial Adverse Reactions	14
8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data	14
8.5 Clinical Trial Adverse Reactions (Adolescents)	16
8.6 Post-Market Adverse Reactions	16

9	DRUG INTERACTIONS.....	17
9.1	Overview	17
9.2	Drug-Drug Interactions.....	17
9.3	Drug-Food Interactions.....	23
9.4	Drug-Herb Interactions.....	23
9.5	Drug-Laboratory Test Interactions	23
10	ACTION AND CLINICAL PHARMACOLOGY.....	23
10.1	Mechanism of Action	23
10.2	Pharmacodynamics.....	23
10.3	Pharmacokinetics.....	24
11	STORAGE, STABILITY AND DISPOSAL.....	28
12	SPECIAL HANDLING INSTRUCTIONS.....	28
PART II: SCIENTIFIC INFORMATION		29
13	PHARMACEUTICAL INFORMATION	29
14	CLINICAL TRIALS	30
14.1	Trial Design and Study Demographics.....	30
14.2	Study Results.....	32
14.3	Comparative Bioavailability Studies.....	38
15	MICROBIOLOGY.....	40
16	NON-CLINICAL TOXICOLOGY.....	42
17	SUPPORTING PRODUCT MONOGRAPHS.....	42
PATIENT MEDICATION INFORMATION.....		43

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

DOVATO (dolutegravir and lamivudine) is indicated as a complete regimen for the treatment of Human Immunodeficiency Virus type 1 (HIV-1) infection in adults and adolescents 12 years of age and older and weighing at least 40 kg.

1.1 Pediatrics

Pediatrics (<18 years of age): Safety and efficacy of DOVATO in pediatric patients less than 12 years of age have not been established. There are no clinical study data with DOVATO in the adolescent population. The safety and efficacy of DOVATO in adolescents 12 years of age and older, and weighing at least 40 kg, is supported by the clinical data from studies of dolutegravir or lamivudine as single agents in combination with other antiretroviral agents in adolescents, and also by the clinical data from studies with dolutegravir in combination with lamivudine in adults (see [CLINICAL TRIALS, Adolescents](#)).

1.2 Geriatrics

Geriatrics (> 65 years of age): Clinical studies of DOVATO did not include sufficient numbers of patients aged 65 years and older to determine whether they respond differently from adult patients < 65 years of age.

2 CONTRAINDICATIONS

DOVATO is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see [DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#).

DOVATO is contraindicated in combination with drugs with narrow therapeutic windows, that are substrates of organic cation transporter 2 (OCT2), including but not limited to dofetilide, and/or fampridine (also known as dalfampridine) (see DRUG INTERACTIONS).

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

- **Post-Treatment Exacerbations of Hepatitis B**
Severe acute exacerbations of hepatitis B have been reported in patients who are infected with hepatitis B virus (HBV) and have discontinued lamivudine, a component of DOVATO. Hepatic function should be monitored closely with both clinical and laboratory follow-up for at least several months in patients who discontinue DOVATO. If appropriate, initiation of anti-hepatitis B therapy may be warranted (see [WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

- As with all antiretroviral drugs, therapy should be initiated by a healthcare professional experienced in the management of HIV infection.
- DOVATO can be taken with or without food.
- DOVATO is a fixed-dose tablet and should not be prescribed for patients requiring dosage adjustments, such as those with creatinine clearance less than 30 mL/min.
- DOVATO is not recommended for patients with any known or suspected viral resistance to dolutegravir or lamivudine.
- DOVATO contains lamivudine and therefore it is recommended to test for Hepatitis B virus (HBV) infection prior to or when initiating DOVATO (see **WARNINGS AND PRECAUTIONS, Hepatic/Biliary/Pancreatic**).
- Perform pregnancy testing before initiation of DOVATO in individuals of childbearing potential.

4.2 Recommended Dose and Dosage Adjustment

The recommended dose of DOVATO in adults and adolescents weighing at least 40 kg is one tablet once daily taken orally.

A separate preparation of dolutegravir (TIVICAY) is available where dose adjustment is required due to drug-drug interactions (see **DRUG INTERACTIONS**).

Pediatrics (< 12 years of age): Safety and efficacy of DOVATO in pediatric patients less than 12 years of age and weighing less than 40 kg have not been established.

Geriatrics (> 65 years of age): Clinical studies of DOVATO did not include sufficient numbers of patients aged 65 years and older to determine whether they respond differently from adult patients < 65 years of age.

Dosage Recommendation with Certain Concomitant Medications

The dolutegravir dose (50 mg) in DOVATO is insufficient when co-administered with medications listed in [Table 1](#) that may decrease dolutegravir concentrations; the following dolutegravir dosage regimen is recommended.

Table 1 Dosing Recommendations for DOVATO with Co-administered Medications

Co-administered Drug	Dosing Recommendation
Oxcarbamazepine, carbamazepine, phenytoin, phenobarbital, St. John's wort or rifampin	Adjust dolutegravir dose to 50 mg twice daily. The additional 50 mg dose of dolutegravir should be taken, separated by 12 hours from DOVATO (see DRUG INTERACTIONS).

Renal insufficiency

DOVATO is not recommended for use in patients with a creatinine clearance less than 30 mL/min as the dose of lamivudine cannot be adjusted (see **ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics**, Special Populations and Conditions, *Renal Insufficiency*, and **WARNINGS AND PRECAUTIONS, Renal**).

Hepatic insufficiency

No dosage adjustment of DOVATO is required in patients with mild or moderate hepatic insufficiency (Child-Pugh score A or B). DOVATO is not recommended in patients with severe hepatic insufficiency (Child-Pugh score C) as it has not been studied in these patients (see **ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics**, Special Populations and Conditions, *Hepatic Insufficiency*).

4.3 Missed Dose

If a dose is missed, patients should take the missed dose as soon as possible unless it is within 4 hours of their next scheduled dose. If a dose is skipped, the patient should not double the next dose.

5 OVERDOSAGE

Symptoms and signs

Experience with overdose of DOVATO or the individual components, dolutegravir and lamivudine is limited. No specific symptoms or signs have been identified.

Treatment

There is no known treatment for overdose with DOVATO. If overdose occurs, the patient should be monitored and standard supportive treatment applied as required. Since lamivudine is dialysable, continuous haemodialysis could be used in the treatment of overdose, although this has not been studied. As dolutegravir is highly bound to plasma proteins, it is unlikely that it will be significantly removed by dialysis.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 2 Dosage Forms, Strengths, Composition and Packaging.

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
oral	tablet 50 mg dolutegravir (as dolutegravir sodium), 300 mg lamivudine	hypromellose, macrogol/PEG, magnesium stearate, mannitol (E421), microcrystalline cellulose, povidone (K29/32), sodium starch glycolate, sodium stearyl fumarate, titanium dioxide

Each film-coated tablet of DOVATO contains 50 mg dolutegravir (as 52.6 mg of dolutegravir sodium) and 300 mg lamivudine.

Dosage Forms

DOVATO tablets are oval, biconvex, white, film-coated tablets, debossed with 'SV 137' on one face.

Packaging

DOVATO tablets are supplied in opaque, white, round, HDPE (high density polyethylene) bottles closed with polypropylene child-resistant closures. Each bottle contains 30 film-coated tablets.

7 WARNINGS AND PRECAUTIONS

Please see the Serious Warnings and Precautions Box at the beginning of Part I: Health Professional Information.

General

DOVATO is a complete regimen for the treatment of HIV-1 infection; therefore, coadministration with other antiretroviral medications for treatment of HIV-1 infection is not recommended.

The safety and efficacy of DOVATO have not been studied in HIV-1-infected patients who have failed previous antiretroviral therapy and are currently not virologically suppressed.

As with other antiretroviral medicinal products, resistance testing and/or historical resistance data should guide the use of DOVATO. DOVATO should not be used in patients with known or suspected resistance to dolutegravir or lamivudine.

Patients receiving DOVATO or any other antiretroviral therapy may still develop opportunistic infections and other complications of HIV infection. Therefore, patients should remain under close clinical observation by physicians experienced in the treatment of these associated HIV diseases.

While effective viral suppression with antiretroviral therapy has been proven to substantially reduce the risk of sexual transmission, a residual risk cannot be excluded. Precautions to prevent transmission should be taken in accordance with national guidelines.

Endocrine and Metabolism**Serum lipids and blood glucose**

Serum lipid and blood glucose levels may increase during antiretroviral therapy. Disease control and life style changes may also be contributing factors. Consideration should be given to the measurement of serum lipids and blood glucose. Lipid disorders and blood glucose elevations should be managed as clinically appropriate.

Hematologic

Very rare occurrences of pure red cell aplasia have been reported with lamivudine use. Discontinuation of lamivudine has resulted in normalization of hematologic parameters in patients with suspected lamivudine-induced pure red cell aplasia.

Hepatic/Biliary/Pancreatic

Hepatotoxicity

Cases of hepatic toxicity including elevated serum liver biochemistries, hepatitis, and acute liver failure have been reported in patients receiving a dolutegravir-containing regimen who had no pre-existing hepatic disease or other identifiable risk factors. Drug-induced liver injury leading to liver transplant has been reported with TRIUMEQ. Monitoring for hepatotoxicity is recommended.

Post-Treatment Exacerbations of Hepatitis B in Patients Co-infected with HIV-1 and HBV

Prior to or when initiating DOVATO, test for HBV infection (see **DOSAGE AND ADMINISTRATION**).

Severe acute exacerbations of HBV have been reported in patients who are co-infected with HBV and HIV-1 and have discontinued lamivudine, a component of DOVATO. Patients who are co-infected with HIV-1 and HBV should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment with DOVATO. If appropriate, initiation of anti-hepatitis B therapy may be warranted, especially in HBV co-infected patients with advanced liver disease or cirrhosis, since post-treatment exacerbation of hepatitis may lead to hepatic decompensation.

Emergence of Lamivudine-Resistant HBV

Emergence of hepatitis B virus variants associated with resistance to lamivudine has also been reported in HIV-1-infected patients who have received lamivudine-containing antiretroviral regimens in the presence of concurrent infection with hepatitis B virus. Consider an alternative regimen in these patients.

Liver chemistry changes in patients with HBV or HCV co-infection

Patients with underlying HBV or HCV may be at increased risk for worsening or development of transaminase elevations with use of a dolutegravir-containing regimen. Liver chemistry elevations consistent with immune reconstitution inflammatory syndrome were observed in some HBV and/or HCV co-infected patients at the start of dolutegravir therapy. Monitoring of liver chemistries is recommended in patients with HBV and/or HCV co-infection. Particular diligence should be applied in initiating or maintaining effective HBV therapy when starting therapy with DOVATO in HBV co-infected patients.

Lactic Acidosis/Severe Hepatomegaly with Steatosis

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues alone or in combination, including lamivudine (a component of DOVATO). Treatment with DOVATO should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity, which may include hepatomegaly and steatosis even in the absence of marked transaminase elevations.

Hypersensitivity Reactions

Hypersensitivity reactions have been reported with integrase inhibitors, including dolutegravir, and were characterized by rash, constitutional findings, and sometimes, organ dysfunction, including liver injury. Discontinue DOVATO and other suspect agents immediately if signs or symptoms of hypersensitivity reactions develop (including, but not limited to, severe rash or rash accompanied by fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions,

conjunctivitis, facial edema, hepatitis, eosinophilia, angioedema). Clinical status including liver aminotransferases should be monitored and appropriate therapy initiated. Delay in stopping treatment with DOVATO or other suspect agents after the onset of hypersensitivity may result in a life-threatening reaction.

Immune

Immune Reconstitution Inflammatory Syndrome

During the initial phase of treatment, patients responding to antiretroviral therapy may develop an inflammatory response to indolent or residual opportunistic infections (such as MAC, CMV, PCP, and TB) which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis and Guillain-Barre syndrome) have also been reported to occur in the setting of immune reconstitution, however the time to onset is more variable, and can occur many months after initiation of treatment and sometimes can be an atypical presentation.

Renal

DOVATO is not recommended for patients with creatinine clearance <30 mL/min because DOVATO is a fixed-dose combination and the dosage of the individual components cannot be adjusted. If a dose reduction of lamivudine, a component of DOVATO, is required for patients with creatinine clearance <30 mL/min, then the individual components should be used (see **DOSAGE AND ADMINISTRATION, Renal Insufficiency**).

Patients with a creatinine clearance between 30 and 49 mL/min receiving DOVATO may experience a 1.6- to 3.3-fold higher lamivudine exposure (AUC) than patients with a creatinine clearance \geq 50 mL/min. There are no safety data from randomized, controlled trials comparing DOVATO to the individual components in patients with a creatinine clearance between 30 and 49 mL/min who received dose-adjusted lamivudine. In the original lamivudine registrational trials in combination with zidovudine, higher lamivudine exposures were associated with higher rates of hematologic toxicities (neutropenia and anemia), although discontinuations due to neutropenia or anemia each occurred in <1% of subjects. Patients with a sustained creatinine clearance between 30 and 49 mL/min who receive DOVATO should be monitored for hematologic toxicities. If new or worsening neutropenia or anemia develop, dose adjustment of lamivudine, per lamivudine prescribing information, is recommended. If lamivudine dose adjustment is indicated, DOVATO should be discontinued and the individual components should be used to construct the treatment regimen.

Sexual Health

Reproduction

Antiretroviral Pregnancy Registry (APR): To monitor maternal-fetal outcomes of pregnant women with HIV exposed to DOVATO and other antiretroviral agents, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients:
<http://www.apregistry.com>
Telephone: (800) 258-4263
Fax: (800) 800-1052

Fertility

There are no data on the effects of dolutegravir or lamivudine on human male or female fertility. Animal studies indicate no effects of dolutegravir or lamivudine on male or female fertility (see [NON-CLINICAL TOXICOLOGY](#)).

7.1 Special Populations

7.1.1 Pregnant Women

DOVATO has not been studied in pregnant women. DOVATO should not be used in pregnant women unless the potential benefits outweigh the potential risks to the fetus. Women of childbearing potential (WOCBP) should be informed about the potential risk of neural tube defects with dolutegravir and counselled about the use of effective contraception. It is recommended that pregnancy testing is conducted prior to initiation of DOVATO. If there are plans to become pregnant, or if pregnancy is confirmed within the first trimester while on DOVATO, the risks and benefits of continuing DOVATO versus switching to another antiretroviral regimen should be discussed with the patient. Factors to consider should include feasibility of switching, tolerability, ability to maintain viral suppression, actual gestational age, risk of transmission to the infant and the available data around the potential risk of neural tube defects and other pregnancy outcomes for dolutegravir and alternative antiretroviral drugs.

In a birth outcome surveillance study in Botswana, a numerically higher rate of neural tube defects was identified with exposure to dolutegravir compared to non-dolutegravir-containing antiretroviral regimens at the time of conception, however, the difference was not statistically significant. Seven cases of neural tube defects were reported in 3,591 deliveries (0.19%) to mothers taking dolutegravir-containing regimens at the time of conception, compared with 21 cases in 19,361 deliveries (0.11%) to mothers taking non-dolutegravir-containing regimens at the time of conception (Prevalence Difference 0.09%; 95% CI -0.031, 0.30). In the same study, an increased risk of neural tube defects was not identified in women who started dolutegravir during pregnancy. Two out of 4,448 deliveries (0.04%) to mothers who started dolutegravir during pregnancy had a neural tube defect, compared with five out of 6,748 deliveries (0.07%) to mothers who started non-dolutegravir-containing regimens during pregnancy. A causal relationship of these events to the use of dolutegravir has not been established. The incidence of neural tube defects in the general population ranges from 0.5-1 case per 1,000 live births. As most neural tube defects occur within the first 4 weeks of fetal development (approximately 6 weeks after the last menstrual period) this potential risk would concern women exposed to dolutegravir at the time of conception and in early pregnancy.

Data analyzed to date from other sources including the Antiretroviral Pregnancy Registry (APR), clinical trials, and post-marketing data are insufficient to address the risk of neural tube defects with dolutegravir. From the APR, one neural tube defect has been identified in 312 (0.32%) live births with periconceptual exposures to dolutegravir.

More than 1,000 outcomes from second and third trimester exposure in pregnant women indicate no evidence of increased risk of adverse birth outcomes and the APR continues to monitor for DTG safety in pregnancy.

Dolutegravir and lamivudine use during pregnancy have been evaluated in the Antiretroviral Pregnancy Registry (APR) in over 700 and 12,800 women, respectively. Available human data from the APR do not show an increased risk of major birth defects for dolutegravir or lamivudine

compared to the background rate.

Based on prospective reports to the APR of over 700 exposures to dolutegravir during pregnancy resulting in live births (including over 450 exposed in the first trimester), there was no difference between the overall risk of birth defects for dolutegravir when compared to the background birth defect rate of 2.7% and 4.17% from two population based surveillance systems (Metropolitan Atlanta Congenital Defects Program (MACDP) with defects of 2.72 per 100 live births and the Texas Birth Defects Registry (TBDR) with 4.17 per 100 live births). The prevalence of defects in live births was 3.5% (95% CI: 2.0% to 5.6%) following first trimester exposure to dolutegravir-containing regimens and 4.2% (95% CI: 2.2% to 7.2%) following second/third trimester exposure to dolutegravir-containing regimens.

Based on prospective reports to the APR of over 12,800 exposures to lamivudine during pregnancy resulting in live births (including over 5300 exposed in the first trimester), there was no difference between the overall risk of birth defects for lamivudine when compared to the background birth defect rate of 2.7% and 4.17% from the MACPD and TBDR, respectively. The prevalence of defects in live births was 3.1% (95% CI: 2.7% to 3.6%) following first trimester exposure to lamivudine-containing regimens and 2.9% (95% CI: 2.5% to 3.3%) following second/third trimester exposure to lamivudine-containing regimens.

In reproductive toxicity studies in animals, dolutegravir was shown to cross the placenta and no evidence of teratogenicity, reproductive function, relevant embryonic or fetal toxicity including neural tube defects was identified in rats and rabbits at ≥ 30 and 0.55 times human clinical exposure based on AUC, respectively.

Lamivudine: There have been reports of mild, transient elevations in serum lactate levels, which may be due to mitochondrial dysfunction, in neonates and infants exposed in utero or peri-partum to nucleoside reverse transcriptase inhibitors (NRTIs). The clinical relevance of transient elevations in serum lactate is unknown. There have also been very rare reports of developmental delay, seizures and other neurological disease. However, a causal relationship between these events and NRTI exposure in utero or peri-partum has not been established. These findings do not affect current recommendations to use antiretroviral therapy in pregnant women to prevent vertical transmission of HIV.

Reproduction studies with lamivudine in rats and rabbits showed no evidence of teratogenicity. Evidence of early embryoletality was seen in the rabbit at lamivudine exposure levels similar to those observed in humans, but there was no indication of this effect in the rat at exposure levels ~ 21 times (based on C_{max}) that of the recommended human dose. Studies in pregnant rats showed that lamivudine is transferred to the fetus through the placenta (see **NON-CLINICAL TOXICOLOGY**).

7.1.2 Breast-feeding

HIV-1-infected mothers should not breast-feed their infants to avoid risking postnatal transmission of HIV. It is expected that dolutegravir will be present in human milk based on animal data. Lamivudine is excreted in human milk at similar concentrations to those found in serum. Because of the potential for HIV-1 transmission and the potential for serious adverse reactions in nursing infants, mothers should be instructed not to breast-feed if they are receiving DOVATO.

7.1.3 Pediatrics

Pediatrics (< 12 years of age): Safety and efficacy of DOVATO have not been established in pediatric patients less than 12 years of age.

7.1.4 Geriatrics

Geriatrics (> 65 years of age): Clinical studies of DOVATO did not include sufficient numbers of patients aged 65 years and older to determine whether they respond differently from adult patients < 65 years of age. In general, caution should be exercised in the administration of DOVATO in elderly patients reflecting the greater frequency of decreased hepatic, renal or cardiac function and of concomitant disease or other drug therapy.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

The following adverse drug reactions are discussed in the WARNINGS AND PRECAUTIONS section:

- Hepatotoxicity
- Severe acute exacerbation of hepatitis B in patients co-infected with HIV-1 and HBV
- Lactic acidosis and severe hepatomegaly with steatosis
- Hypersensitivity reactions
- Immune reconstitution inflammatory syndrome

8.2 Clinical Trial Adverse Reactions

Because clinical trials are conducted under very specific conditions, the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

For details on adverse reactions that have occurred in studies with dolutegravir or lamivudine, please refer to the TIVICAY and 3TC product monographs.

Treatment-Naïve Patients

The safety assessment of DOVATO in HIV-1-infected, treatment naïve adult patients with viral load \leq 500,000 HIV-1 RNA copies per mL is based on the pooled primary Week 48 and Week 96 analyses of data from two identical, multicenter, double-blind, controlled trials, GEMINI-1 and GEMINI-2 where dolutegravir plus lamivudine were co-administered as single agents (TIVICAY and 3TC).

A total of 1,433 adult HIV-1-infected treatment-naïve subjects were randomized to dolutegravir 50 mg plus lamivudine 300 mg, as a complete regimen once daily, or dolutegravir 50 mg plus fixed-dose combination tenofovir disoproxil fumarate (TDF)/emtricitabine (FTC), administered once daily.

The rates of adverse events leading to discontinuation in the pooled analysis were 2% (Week 48) and 3% (Week 96) of subjects in both treatment arms. The most common adverse events leading to discontinuation were psychiatric disorders (<1% (Week 48) and 1% (Week 96) of subjects in both treatment arms).

Adverse reactions (all grades) observed in at least 2% of subjects in either treatment arm of either individual study or the pooled analysis of GEMINI-1 and GEMINI-2 studies at Weeks 48 and 96 are provided in [Table 3](#).

The adverse reactions observed for TIVICAY plus 3TC in the pooled Week 48 and Week 96 analyses from GEMINI-1 and GEMINI-2 were generally consistent with the adverse reaction profiles and severities for the individual components, when administered with other antiretroviral agents. The majority of adverse reactions related to TIVICAY plus 3TC were of Grade 1 intensity.

Table 3 Treatment-Emergent Adverse Reactions (All Grades) and at Least 2% Frequency in either treatment arm (Week 48 and 96 Individual Studies and Pooled Analyses)

System Organ Class/ Preferred Term	GEMINI-1				GEMINI-2				POOLED			
	TIVICAY + 3TC (n = 356) n (%)		TIVICAY + TRUVADA (n=358) n (%)		TIVICAY + 3TC (n = 360) n (%)		TIVICAY + TRUVADA (n=359) n (%)		TIVICAY + 3TC (n = 716) n (%)		TIVICAY + TRUVADA (n = 717) n (%)	
	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96
Nervous system disorders												
Headache	14 (4)	14 (4)	19 (5)	19 (5)	7 (2)	7 (2)	11 (3)	11 (3)	21 (3)	21 (3)	30 (4)	30 (4)
Somnolence	7 (2)	7 (2)	5 (1)	5 (1)	1 (<1)	1 (<1)	2 (<1)	2 (<1)	8 (1)	8 (1)	7 (<1)	7 (<1)
Dizziness	5 (1)	5 (1)	6 (2)	6 (2)	3 (<1)	3 (<1)	7 (2)	7 (2)	8 (1)	8 (1)	13 (2)	13 (2)
Gastrointestinal disorders												
Nausea	8 (2)	8 (2)	23 (6)	23 (6)	6 (2)	6 (2)	16 (4)	16 (4)	14 (2)	14 (2)	39 (5)	39 (5)
Diarrhea	6 (2)	7 (2)	14 (4)	14 (4)	8 (2)	8 (2)	5 (1)	5 (1)	14 (2)	15 (2)	19 (3)	19 (3)
Psychiatric disorders												
Insomnia	7 (2)	8 (2)	12 (3)	12 (3)	6 (2)	7 (2)	6 (2)	7 (2)	13 (2)	15 (2)	18 (3)	19 (3)
Anxiety	N/A	5 (1)	N/A	1 (<1)	N/A	6 (2)	N/A	4 (1)	N/A	11 (2)	N/A	5 (<1)
General disorders and administration site conditions												
Fatigue	6 (2)	7 (2)	3 (<1)	3 (<1)	4 (1)	4 (1)	3 (<1)	3 (<1)	10 (1)	11 (2)	6 (<1)	6 (<1)

The only adverse reaction of \geq Grade 2 occurring in $\geq 1\%$ of subjects treated with TIVICAY plus 3TC was headache (1%).

Virologically Suppressed Patients

The safety of DOVATO in virologically suppressed adults was based on Week 48 data from 740 subjects in a randomized, parallel-group, open-label, multicenter, non-inferiority controlled trial (TANGO). Subjects who were on a stable suppressive tenofovir alafenamide-based regimen (TBR) were randomized to receive DOVATO once daily or continue with TBR. Overall, the

safety profile of DOVATO in virologically suppressed adult subjects in the TANGO trial was similar to that of TIVICAY plus 3TC in treatment naïve subjects in the GEMINI trials.

8.3 Less Common Clinical Trial Adverse Reactions

The following adverse reactions occurred in less than 2% of patients receiving dolutegravir plus lamivudine or are from studies described in the product monographs of the individual components TIVICAY (dolutegravir) and 3TC (lamivudine). Some events have been included because of their seriousness and assessment of potential causal relationship.

Blood and Lymphatic Systems Disorders: Anemia, neutropenia, thrombocytopenia.

Gastrointestinal Disorders: Abdominal pain, abdominal discomfort, flatulence, upper abdominal pain, vomiting

General Disorders: Fever, malaise

Hepatobiliary Disorders: Hepatitis

Immune System Disorders: Hypersensitivity, immune reconstitution inflammatory syndrome

Musculoskeletal and Connective Tissue Disorders: Myositis

Psychiatric Disorders: Abnormal dreams, depression, suicidal ideation or suicide attempt (particularly in patients with a pre-existing history of depression or psychiatric illness)

Renal and Urinary Disorders: Renal impairment

Skin and Subcutaneous Tissue Disorders: Pruritus, rash

8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data

Selected laboratory abnormalities with a worsening grade from baseline and representing the worst-grade toxicity in GEMINI-1 and GEMINI 2 are presented in [Table 4](#).

Table 4 Selected Laboratory Abnormalities (Grades 2 and 3 to 4; Week 48 and 96 Pooled Analyses) in GEMINI-1 and GEMINI-2 Trials

Laboratory Parameter Preferred Term	Week 48		Week 96	
	TIVICAY + 3TC (N = 716) n (%)	TIVICAY + TRUVADA (N = 717) n (%)	TIVICAY + 3TC (N = 716) n (%)	TIVICAY + TRUVADA (N = 717) n (%)
ALT				
Grade 2 (>2.5 to 5.0 x ULN)	13 (2)	20 (3)	18 (3)	27 (4)
Grade 3 to 4 (>5.0 x ULN)	18 (3)	18 (3)	23 (3)	19 (3)
AST				
Grade 2 (>2.5 to 5.0 x ULN)	22 (3)	19 (3)	29 (4)	27 (4)
Grade 3 to 4 (>5.0 x ULN)	12 (2)	24 (3)	18 (3)	25 (3)
Total Bilirubin				
Grade 2 (1.6 to 2.5 x ULN)	9 (1)	17 (2)	16 (2)	23 (3)
Grade 3 to 4 (>2.5 x ULN)	7 (<1)	7 (<1)	8 (1)	7 (<1)
Cholesterol				
Grade 2 (6.19 to <7.77 mmol/L)	30 (4)	14 (2)	37 (5)	18 (3)
Grade 3 to 4 (>7.77 mmol/L)	0	0	0	1 (<1)
Creatine kinase				
Grade 2 (6.0 to 9.9 x ULN)	26 (4)	21 (3)	29 (4)	31 (4)
Grade 3 to 4 (≥10.0 x ULN)	32 (4)	35 (5)	46 (6)	47 (7)
Hyperglycemia				
Grade 2 (6.95 to 13.89 mmol/L)	48 (7)	29 (4)	62 (9)	46 (6)
Grade 3 to 4 (>13.89 mmol/L)	5 (<1)	5 (<1)	6 (<1)	5 (<1)
LDL Cholesterol				
Grade 2 (4.12 to < 4.9 mmol/L)	20 (3)	12 (2)	25 (3)	15 (2)
Grade 3 to 4 (> 4.9 mmol/L)	8 (1)	3 (<1)	8 (1)	5 (<1)
Lipase				
Grade 2 (>1.5 to 3.0 x ULN)	37 (5)	34 (5)	41 (6)	45 (6)
Grade 3 to 4 (>3.0 x ULN)	7 (<1)	19 (3)	15 (2)	29 (4)
Triglycerides				
Grade 2 (>3.42 to 5.7 mmol/L)	13 (2)	13 (2)	18 (3)	18 (3)
Grade 3 to 4 (> 5.7 mmol/L)	9 (1)	4 (<1)	10 (1)	4 (<1)

ULN = Upper limit of normal, ALT = Alanine Aminotransferase, AST = Aspartate Aminotransferase

Changes in Clinical Laboratory Values

Dolutegravir has been shown to increase serum creatinine due to inhibition of tubular secretion of creatinine without affecting renal glomerular function. Increases in serum creatinine occurred within the first four weeks of treatment with dolutegravir plus lamivudine and remained stable through 96 weeks. A mean change from baseline of 12.25 µmol/L (range: 34.5 µmol/L to 99.9 µmol/L) was observed after 96 weeks of treatment (see [ACTION AND CLINICAL PHARMACOLOGY](#), [Pharmacodynamics](#), Effects on Renal Function).

Small increases in total bilirubin (without clinical jaundice) were observed with dolutegravir plus lamivudine. These changes are not considered clinically relevant as they likely reflect competition between dolutegravir and unconjugated bilirubin for a common clearance pathway (UGT1A1) (see **ACTION AND CLINICAL PHARMACOLOGY, Pharmacokinetics, Metabolism**).

In the pooled analysis, few subjects in either group experienced changes in lipid profile. A small number of subjects in each treatment group experienced LDL cholesterol toxicities \geq Grade 2, TIVICAY + 3TC 4% (Week 48) and 5% (Week 96) and TIVICAY + TRUVADA group 2% (Week 48) and 3% (Week 96). However, both treatment groups showed an overall reduction in the mean total cholesterol/HDL ratio, with a greater reduction in the TIVICAY + TRUVADA group. A small proportion of subjects in both treatment groups also had emergent triglyceride toxicities of \geq Grade 2, TIVICAY + 3TC 3% (Week 48) and 4% (Week 96), TIVICAY + TRUVADA 2% (Week 48) and 3% (Week 96). A total of 40 and 16 subjects receiving TIVICAY + 3TC and TIVICAY + TRUVADA, respectively, initiated lipid-lowering agents post-baseline by Week 96.

Table 5 Mean Change from Baseline in Fasted Lipid Values (Week 48 and 96 Pooled Analyses) in GEMINI-1 and GEMINI-2 Trials

Laboratory Parameter Preferred Term	Week 48		Week 96	
	TIVICAY plus 3TC (n = 716)	TIVICAY plus TRUVADA (n = 717)	TIVICAY plus 3TC (n = 716)	TIVICAY plus TRUVADA (n = 717)
Cholesterol (mmol/L)	0.35	-0.18	0.39	-0.14
HDL cholesterol (mmol/L)	0.15	0.02	0.19	0.08
LDL cholesterol (mmol/L)	0.19	-0.16	0.16	-0.17
Triglycerides (mmol/L)	0.04	-0.08	0.13	-0.12
Total cholesterol/HDL cholesterol ratio	-0.09	-0.26	-0.13	-0.42

8.5 Clinical Trial Adverse Reactions (Adolescents)

There are no clinical study data with DOVATO in the adolescent population. However, a summary of the clinical trial adverse reactions from prior adolescent studies of dolutegravir or lamivudine are available in the respective TIVICAY, 3TC and TRIUMEQ product monographs. For the adolescent studies, there were no additional types of adverse reactions beyond those observed in the adult population.

8.6 Post-Market Adverse Reactions

These events have been chosen for inclusion due to either their seriousness, frequency of reporting, potential causal connection to dolutegravir- and/or lamivudine-containing regimens, or a combination of these factors. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made.

Body as a whole: anaphylaxis, weakness

Blood and Lymphatic Systems Disorders: pure red cell aplasia, anemia, lymphadenopathy

Gastrointestinal Disorders: rises in serum amylase, pancreatitis, stomatitis

Hepatobiliary Disorders: acute hepatic failure, splenomegaly

Investigations: weight increased

Metabolism and Nutrition Disorders: lactic acidosis, hyperlactatemia, hepatic steatosis, hyperglycemia

Musculoskeletal and connective tissue disorders: muscle disorders including rarely rhabdomyolysis, arthralgia, myalgia

Nervous System Disorders: paresthesia, peripheral neuropathy

Skin and Subcutaneous Tissue Disorders: alopecia, urticaria, pruritus

9 DRUG INTERACTIONS

9.1 Overview

DOVATO contains dolutegravir plus lamivudine and any interactions that have been identified with either component individually may occur with DOVATO. There are no significant interactions between dolutegravir and lamivudine. Because DOVATO is a complete regimen coadministration with other antiretroviral medications for the treatment of HIV-1 infection is not recommended and information regarding potential drug-drug interactions with other antiretroviral medications is not provided. For more information on these interactions, please refer to the TIVICAY and 3TC product monographs.

9.2 Drug-Drug Interactions

Effect of Dolutegravir or Lamivudine on the Pharmacokinetics of Other Agents

Dolutegravir

In vitro, dolutegravir did not inhibit ($IC_{50} > 50 \mu M$) the enzymes: cytochrome P450 (CYP)1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A, uridine diphosphate glucuronosyl transferase (UGT)1A1 or UGT2B7, or transporters: P-glycoprotein (Pgp), breast cancer resistance protein (BCRP), bile salt export pump (BSEP), organic anion transporter polypeptide (OATP)1B1, OATP1B3, OCT1, or multidrug resistance protein (MRP)2 or MRP4. *In vitro*, dolutegravir did not induce CYP1A2, CYP2B6, or CYP3A4. Based on these data, dolutegravir is not expected to affect the pharmacokinetics of drugs that are substrates of these enzymes or transporters.

In vitro, dolutegravir inhibited the renal organic cation transporter 2, OCT2 ($IC_{50} = 1.93 \mu M$), multidrug and toxin extrusion transporter (MATE) 1 ($IC_{50} = 6.34 \mu M$) and MATE2-K ($IC_{50} = 24.8 \mu M$). *In vivo*, dolutegravir has a low potential to affect the transport of MATE2-K substrates. *In vivo*, dolutegravir inhibits tubular secretion of creatinine by inhibiting OCT2. Dolutegravir may increase plasma concentrations of drugs in which excretion is dependent upon OCT2 (for example dofetilide, fampridine (also known as dalfampridine) (see [CONTRAINDICATIONS](#)), metformin) or MATE1 (see [Table 6](#)).

In vitro, dolutegravir inhibited the basolateral renal transporters: organic anion transporter (OAT) 1 ($IC_{50} = 2.12 \mu M$) and OAT3 ($IC_{50} = 1.97 \mu M$). Based upon the dolutegravir unbound plasma concentration, *in silico* modelling, and no notable effect on the pharmacokinetics *in vivo* of the

OAT substrates tenofovir and para-aminohippurate, dolutegravir has low propensity to cause drug interactions via inhibition of OAT transporters.

Lamivudine

In vitro, lamivudine does not inhibit or induce CYP enzymes (such as CYP 3A4, CYP 2C9 or CYP 2D6). Lamivudine demonstrates no or weak inhibition of the drug transporters organic anion transporter 1B1 (OATP1B1), OATP1B3, breast cancer resistance protein (BCRP) or P-glycoprotein (Pgp), multidrug and toxin extrusion protein 1 (MATE1), MATE2-K or organic cation transporter 3 (OCT3). Lamivudine is therefore not expected to affect the plasma concentrations of drugs that are substrates of these enzymes or drug transporters.

Lamivudine is an inhibitor of OCT1 and OCT2 *in vitro* with IC50 values of 17 and 33 μ M, respectively, however lamivudine has low potential to affect the plasma concentrations of OCT1 and OCT2 substrates at therapeutic drug exposures (up to 300 mg).

Effect of Other Agents on the Pharmacokinetics of Dolutegravir or Lamivudine

Dolutegravir

Dolutegravir is metabolized by UGT1A1 with some contribution from CYP3A. Dolutegravir is also a substrate of UGT1A3, UGT1A9, Pgp and BCRP *in vitro*; therefore drugs that induce those enzymes and transporters, may decrease dolutegravir plasma concentration and reduce the therapeutic effect of dolutegravir.

Co-administration of dolutegravir and other drugs that inhibit UGT1A1, UGT1A3, UGT1A9, CYP3A4 and/or Pgp may increase dolutegravir plasma concentration (see [Table 6](#)).

In vitro, dolutegravir is not a substrate of human organic anion transporting polypeptide (OATP)1B1, OATP1B3, or OCT1.

Coadministration of dolutegravir with polyvalent cation-containing products may lead to decreased adsorption of dolutegravir.

Lamivudine

Lamivudine is a substrate of MATE1, MATE2-K and OCT2 *in vitro*. Trimethoprim (an inhibitor of these drug transporters) has been shown to increase lamivudine plasma concentrations, however this interaction is not considered clinically significant as no dose adjustment of lamivudine is needed.

Lamivudine is a substrate of the hepatic uptake transporter OCT1. As hepatic elimination plays a minor role in the clearance of lamivudine, drug interactions due to inhibition of OCT1 are unlikely to be of clinical significance.

Lamivudine is a substrate of Pgp and BCRP, however due to its high bioavailability it is unlikely that these transporters play a significant role in the absorption of lamivudine. Therefore, co-administration of drugs that are inhibitors of these efflux transporters is unlikely to affect the disposition and elimination of lamivudine.

Lamivudine is predominantly eliminated by active organic cationic secretion. The possibility of interactions with other drugs administered concurrently should be considered, particularly when the main route of administration is renal.

Established or Potential Drug Interactions

Established and theoretical interactions with selected medicinal products are listed in [Table 6](#). The drugs listed in this table are not all-inclusive. Recommendations are based on either drug interaction studies, or potential or predicted interactions due to the expected magnitude of interaction and/or potential for serious adverse events or loss of efficacy.

Table 6 Established or Potential Drug-Drug Interactions

Concomitant Drug Class: Drug Name	Effect on Concentration of Dolutegravir, Lamivudine, or Concomitant Drug*	Clinical Comment
DOLUTEGRAVIR		
Antiarrhythmic: Dofetilide	Dofetilide↑	Coadministration of DOVATO with dofetilide is contraindicated due to potential life-threatening toxicity caused by high dofetilide concentrations.
Potassium channel blocker: Fampridine (also known as dalfampridine)	Fampridine/dalfampridine↑	Coadministration is contraindicated with DOVATO due to the potential for seizures associated with fampridine/dalfampridine.
Anticonvulsants Oxcarbazepine Phenytoin Phenobarbital Carbamazepine ^a	Dolutegravir↓	An additional 50 mg dose of dolutegravir (TIVICAY) should be taken, separated by 12 hours from DOVATO.
Antidiabetics: Metformin ^a	Co-administered with DOVATO: Metformin ↑	Consider metformin dose adjustments when starting or stopping concomitant treatment to maintain glycemic control.
Antimycobacterials Rifampin ^a	Dolutegravir↓	An additional 50 mg dose of dolutegravir (TIVICAY) should be taken, separated by 12 hours from DOVATO.
Medications containing polyvalent cations (e.g. Mg, Al) Cation-containing antacids ^a or laxative, sucralfate, buffered medications	Dolutegravir↓	DOVATO is recommended to be administered 2 hours before or 6 hours after taking medications containing polyvalent cations.
Calcium and iron supplements ^a	Dolutegravir ↓	When taken with food,

Concomitant Drug Class: Drug Name	Effect on Concentration of Dolutegravir, Lamivudine, or Concomitant Drug*	Clinical Comment
Includes multivitamins that contain calcium or iron.		DOVATO and calcium and/or iron supplements or multivitamins containing calcium and/or iron can be taken at the same time. Under fasting conditions, DOVATO should be taken 2 hours before or 6 hours after taking supplements containing calcium and/or iron.
LAMIVUDINE		
Trimethoprim/sulfamethoxazole (Co-trimoxazole)	Lamivudine: AUC ↑ ~40% Trimethoprim: AUC ↔ Sulfamethoxazole: AUC ↔	Unless the patient has renal impairment, no dosage adjustment of DOVATO is necessary. Co-administration with DOVATO is not recommended in patients with renal impairment as lamivudine dosage adjustment is not possible. The effect of coadministration of lamivudine with higher doses of co-trimoxazole used for the treatment of <i>Pneumocystis jiroveci</i> pneumonia (often referred to as PCP) and toxoplasmosis has not been studied.
Sorbitol solution (3.2 g, 10.2 g, 13.4 g)	Single dose lamivudine oral solution 300 mg Lamivudine: AUC ↓ 14%; 32%; 36% C _{max} ↓ 28%; 52%, 55%.	When possible, avoid chronic coadministration of sorbitol-containing medicines with DOVATO. Consider more frequent monitoring of HIV-1 viral load when chronic coadministration cannot be avoided.

↑ = Increase, ↓ = Decrease, ↔ = No change.

^a For magnitude of interaction, see [Table 7](#) and [Table 8](#).

The effects of DTG on the exposure of co-administered drugs are shown in [Table 7](#). The effects of co-administered drugs on the exposure of DTG are shown in [Table 8](#).

Table 7 Summary of Effect of Dolutegravir on the Pharmacokinetics of Co-administered Drugs

Co-administered Drug(s) and Dose(s)	Dose of Dolutegravir	n	Geometric Mean Ratio (90% CI) of Pharmacokinetic Parameters of Co-administered Drug With/Without Dolutegravir No Effect = 1.00		
			C _r or C ₂₄	AUC	C _{max}
Daclatasvir 60 mg once daily	50 mg once daily	12	1.06 (0.88 to 1.29)	0.98 (0.83 to 1.15)	1.03 (0.84 to 1.25)
Ethinyl estradiol 0.035 mg	50 mg twice daily	15	1.02 (0.93, 1.11)	1.03 (0.96, 1.11)	0.99 (0.91, 1.08)
Methadone 16 to 150 mg	50 mg twice daily	12	0.99 (0.91, 1.07)	0.98 (0.91, 1.06)	1.00 (0.94, 1.06)
Midazolam 3 mg	25 mg once daily	10	–	0.95 (0.79, 1.15)	–
Norgestimate 0.25 mg	50 mg twice daily	15	0.93 (0.85, 1.03)	0.98 (0.91, 1.04)	0.89 (0.82, 0.97)
Metformin 500 mg twice daily	50 mg once daily	14	–	1.79 (1.65, 1.93)	1.66 (1.53, 1.81)
Metformin 500 mg twice daily	50 mg twice daily	14	–	2.45 (2.25, 2.66)	2.11 (1.91, 2.33)

Table 8 Summary of Effect of Co-administered Drugs on the Pharmacokinetics of Dolutegravir

Co-administered Drug(s) and Dose(s)	Dose of Dolutegravir	n	Geometric Mean Ratio (90% CI) of Dolutegravir Pharmacokinetic Parameters With/Without Co-administered Drugs No Effect = 1.00		
			C _r or C ₂₄	AUC	C _{max}
Maalox®	50 mg single dose	16	0.26 (0.21, 0.31)	0.26 (0.22, 0.32)	0.28 (0.23, 0.33)
Maalox® 2 hrs after dolutegravir	50 mg single dose	16	0.70 (0.58, 0.85)	0.74 (0.62, 0.90)	0.82 (0.69, 0.98)
Calcium Carbonate 1200 mg simultaneous administration (fasted)	50 mg single dose	12	0.61 (0.47, 0.80)	0.61 (0.47, 0.80)	0.63 (0.50, 0.81)
Calcium Carbonate 1200 mg simultaneous administration (fed)	50 mg single dose	11	1.08 (0.81, 1.42)	1.09 (0.84, 1.43)	1.07 (0.83, 1.38)
Calcium Carbonate 1200 mg 2 hrs after dolutegravir	50 mg single dose	11	0.90 (0.68, 1.19)	0.94 (0.72, 1.23)	1.00 (0.78, 1.29)
Ferrous Fumarate 324 mg simultaneous administration (fasted)	50 mg single dose	11	0.44 (0.36, 0.54)	0.46 (0.38, 0.56)	0.43 (0.35, 0.52)
Ferrous Fumarate 324 mg simultaneous administration (fed)	50 mg single dose	11	1.00 (0.81, 1.23)	0.98 (0.81, 1.20)	1.03 (0.84, 1.26)
Ferrous Fumarate 324 mg 2 hrs after dolutegravir	50 mg single dose	10	0.92 (0.74, 1.13)	0.95 (0.77, 1.15)	0.99 (0.81, 1.21)
Multivitamin One tablet once daily	50 mg single dose	16	0.68 (0.56, 0.82)	0.67 (0.55, 0.81)	0.65 (0.54, 0.77)
Omeprazole 40 mg once daily	50 mg single dose	12	0.95 (0.75, 1.21)	0.97 (0.78, 1.20)	0.92 (0.75, 1.11)
Prednisone 60 mg once daily with taper	50 mg once daily	12	1.17 (1.06, 1.28)	1.11 (1.03, 1.20)	1.06 (0.99, 1.14)
Rifampin ^a 600 mg once daily	50 mg twice daily ^a	11	0.28 (0.23, 0.34)	0.46 (0.38, 0.55)	0.57 (0.49, 0.65)
Rifampin ^b 600 mg once daily	50 mg twice daily ^b	11	1.22 (1.01, 1.48)	1.33 (1.15, 1.53)	1.18 (1.03, 1.37)
Rifabutin 300 mg once daily	50 mg once daily	9	0.70 (0.57, 0.87)	0.95 (0.82, 1.10)	1.16 (0.98, 1.37)
Carbamazepine 300 mg twice daily	50 mg once daily	14	0.27 (0.24, 0.31)	0.51 (0.48, 0.55)	0.67 (0.61, 0.73)
Daclatasvir 60 mg once daily	50 mg once daily	12	1.45 (1.25 to 1.68)	1.33 (1.11 to 1.59)	1.29 (1.07 to 1.57)

^a Comparison is rifampin taken with dolutegravir 50 mg twice daily compared with dolutegravir 50 mg twice daily.

^b Comparison is rifampin taken with dolutegravir 50 mg twice daily compared with dolutegravir 50 mg once daily.

Drugs with No Observed or Predicted Interactions with DOVATO

Based on drug interaction studies conducted with DOVATO or the components of DOVATO, no clinically significant drug interactions have been either observed or are expected when DOVATO is administered with the following drugs: hormonal contraceptives containing norgestimate and ethinyl estradiol, methadone, midazolam, omeprazole, prednisone, rifabutin, daclatasvir, sofosbuvir/velpatasvir, trimethoprim-sulfamethoxazole (except in renal impairment, see [Table 6](#)), and calcium carbonate, ferrous fumarate, or cation-containing multivitamin supplements (when taken with food, see [Table 6](#)).

9.3 Drug-Food Interactions

DOVATO can be taken with or without food (see [ACTION AND CLINICAL PHARMACOLOGY](#)).

9.4 Drug-Herb Interactions

No interaction study has been conducted, however, St. John's Wort is a potent CYP3A inducer and may potentially decrease dolutegravir plasma concentration. In adults and adolescent patients, an additional dose of TIVICAY 50 mg separated by 12 hours from DOVATO may be considered when taken together with St. John's Wort.

9.5 Drug-Laboratory Test Interactions

No Drug-Laboratory interactions have been identified.

10 ACTION AND CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Dolutegravir inhibits HIV integrase by binding to the integrase active site and blocking the strand transfer step of retroviral Deoxyribonucleic acid (DNA) integration which is essential for the HIV replication cycle. *In vitro*, dolutegravir dissociates slowly from the active site of the wild type integrase-DNA complex ($t_{1/2}$ 71 hours). Strand transfer biochemical assays using purified HIV-1 integrase and pre-processed substrate DNA resulted in IC_{50} values of 2.7 nM and 12.6 nM.

Lamivudine is a synthetic nucleoside analogue, an (-) enantiomer of a dideoxy analogue of cytidine. Lamivudine is metabolized by intracellular kinases to its triphosphate (TP), which is the active moiety (lamivudine triphosphate or L-TP). Lamivudine is a nucleoside reverse transcriptase inhibitor (NRTI), and is a potent, selective inhibitor of HIV-1 and HIV-2 replication *in vitro*. *In vitro* L-TP has an intracellular half-life of approximately 10.5 to 15.5 hours. L-TP is a substrate for and a competitive inhibitor of HIV reverse transcriptase (RT). Inhibition of RT is via viral DNA chain termination after nucleoside analogue incorporation. L-TP shows significantly less affinity for host cell DNA polymerases and is a weak inhibitor of mammalian α , β , and γ DNA polymerases.

10.2 Pharmacodynamics

In a randomized, dose-ranging trial, HIV 1 infected patients treated with dolutegravir monotherapy demonstrated rapid and dose-dependent antiviral activity with mean declines from

baseline to Day 11 in HIV-1 RNA of 1.5, 2.0, and 2.5 log₁₀ for dolutegravir 2 mg, 10 mg, and 50 mg once daily, respectively. This antiviral response was maintained for 3 to 4 days after the last dose in the 50 mg group.

Effects on Electrocardiogram

Dolutegravir

In a randomized, placebo-controlled, cross-over trial, 42 healthy subjects received single dose oral administrations of placebo, dolutegravir 250 mg suspension (exposures approximately 3-fold of the 50 mg once-daily dose at steady state), and moxifloxacin (400 mg, active control) in random sequence. Dolutegravir did not prolong the QTc interval for 24 hours post dose. After baseline and placebo adjustment, the maximum mean QTc change based on Fridericia correction method (QTcF) was 1.99 msec (1-sided 95% upper CI: 4.53 msec).

Effects on Renal Function

The effect of dolutegravir on serum creatinine clearance (CrCl), glomerular filtration rate (GFR) using iohexol as the probe and effective renal plasma flow (ERPF) using para-aminohippurate (PAH) as the probe was evaluated in an open-label, randomized, 3-arm, parallel, placebo-controlled study in 37 healthy subjects, who were administered dolutegravir 50 mg once daily (n=12), 50 mg twice daily (n=13) or placebo once daily (n=12) for 14 days. A modest decrease in CrCl was observed with dolutegravir within the first week of treatment, consistent with that seen in clinical studies. Dolutegravir at both doses had no significant effect on GFR or ERPF. These data support *in vitro* studies which suggest that the small increases in creatinine observed in clinical studies are due to the nonpathologic inhibition of the organic cation transporter 2 (OCT2) in the proximal renal tubules, which mediates the tubular secretion of creatinine.

In the pooled analysis of GEMINI-1 and GEMINI-2 studies in treatment-naïve adult patients at the Week 48 and 96 analyses, TIVICAY + 3TC had an increase in the estimated GFR using cystatin C adjusted CKD-EPI equation (adjusted mean change from baseline of 6.3 (Week 48) and 10.7 (Week 96) mL/min/1.73 m²). Change from baseline analysis showed that urine albumin/creatinine and protein/creatinine ratios decreased at Week 48 and 96 compared to baseline in the TIVICAY + 3TC group (urine albumin/creatinine Week 48/baseline ratio of 0.914 and urine albumin/creatinine Week 96/baseline ratio of 0.934; protein/creatinine Week 48/baseline ratio of 0.869 and protein/creatinine Week 96/baseline ratio of 0.878).

10.3 Pharmacokinetics

In a fasted comparative bioavailability study, the dolutegravir C_{max} was equivalent and the dolutegravir AUC_T was 16% higher when comparing the DOVATO tablet to dolutegravir 50 mg co-administered with lamivudine 300 mg. The higher DTG AUC_T does not significantly affect patient safety or antiviral efficacy based on historical clinical efficacy and safety data for DTG 50 mg BID. The lamivudine AUC_T was equivalent when comparing the DOVATO tablet to lamivudine 300 mg co-administered with dolutegravir 50 mg. Lamivudine C_{max} for the DOVATO tablet was 32% higher than lamivudine 300 mg co-administered with dolutegravir 50 mg. The higher lamivudine C_{max}, which reflects differences in the rate of absorption but not extent of absorption, does not significantly affect patient safety or antiviral efficacy based on historical clinical efficacy and safety data at higher lamivudine doses/exposures. Following multiple oral doses of DOVATO in HIV-infected, treatment experienced subjects in the Phase III TANGO

study, the steady state dolutegravir and lamivudine AUC and C_{max} were similar to historical exposures.

The pharmacokinetic (PK) properties of the components of DOVATO are provided in [Table 9](#).

Table 9 Pharmacokinetic Properties of the Components of DOVATO

	Dolutegravir	Lamivudine
Absorption		
AUC _T ^a (μg.h/mL)	52.3 (31.5)	13.4 (18.1)
C _{max} ^a (μg/mL)	2.91 (30.6)	3.22 (29.3)
T _{max} ^a (h)	2.5 (0.5, 6.0)	1 (0.5, 3.5)
Effect of high-fat meal (relative to fasting) ^b	AUC _i % Ratio 132.6 (118.4, 148.5)	AUC _i % Ratio 91.1 (86.6, 95.9)
Distribution		
% Bound to human plasma proteins	~99	<36
Source of protein binding data	<i>in vitro</i>	<i>in vitro</i>
Blood-to-plasma ratio	0.44-0.54	1.1 - 1.2 ^c
Metabolism		
Metabolic pathways	UGT1A1 CYP3A (minor)	Not significantly metabolized
Elimination		
Major route of elimination	Metabolism	Renal, by the organic cationic transport system
t _{1/2} (h)	~14	18-19
% of dose excreted as total [¹⁴ C] (unchanged drug) in urine	31 (<1) ^d	ND (~70) ^e
% of dose excreted as total [¹⁴ C] (unchanged drug) in feces	64 (53) ^d	ND (ND)

ND: not determined

- Single dose PK parameters presented as geometric mean (CVb) except for T_{max} which is presented as median (range)
- Geometric mean ratio (fed/fasted) in PK parameters and (90% confidence interval). High-calorie/high-fat meal = ~900 kcal, 56% fat
- Lamivudine blood-to-plasma ratio (B/P) was calculated based on the percent (p) of blood lamivudine associated with erythrocytes (53% to 57%) and the hematocrit value (H) using the equation B/P = (1-H)(1-p)
- Based on single-dose, mass-balance study of [¹⁴C] dolutegravir
- Based on 24-hour urine collection obtained after oral or IV administration (NUCB1001)

Absorption: Dolutegravir and lamivudine are rapidly absorbed following oral administration. The absolute bioavailability of dolutegravir has not been established. The absolute bioavailability of oral lamivudine in adults is 80 to 85%. For DOVATO, the median time to maximal plasma concentrations (t_{max}) is 2.5 hours for dolutegravir and 1.0 hour for lamivudine, when dosed under fasted conditions.

Following multiple oral doses of dolutegravir 50 mg once daily, the geometric mean steady state pharmacokinetic parameter estimates are 53.6 μg.h/mL for AUC₂₄, 3.67 μg/mL for C_{max}, and 1.11 μg/mL for C₂₄. Following multiple-dose oral administration of lamivudine 300 mg once daily for seven days the mean steady-state C_{max} is 2.04 μg/mL and the mean AUC₂₄ is 8.87 μg.h/mL.

Effect of Food on Oral Absorption

DOVATO may be administered with or without food. Administration of DOVATO with a high-fat, high-calorie meal increased dolutegravir AUC_T and C_{max} by 32% and 21%, respectively, and decreased the lamivudine C_{max} by 32% compared to fasted conditions. The lamivudine AUC_T was not affected by a high-fat, high-calorie meal. These changes are not clinically significant.

Distribution: Dolutegravir is highly bound ($\geq 98.9\%$) to human plasma proteins based on *in vivo* data and binding is independent of plasma dolutegravir concentration. The apparent volume of distribution (Vd/F) following 50 mg once daily oral administration was estimated at 17.4 L based on population pharmacokinetic analysis. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays low plasma protein binding (less than 36%).

Cerebrospinal Fluid (CSF)

In 12 treatment-naïve patients on dolutegravir plus abacavir/lamivudine, the median dolutegravir concentration in CSF was 18 ng/mL (ranging from 4 to 23 ng/mL) 2 to 6 hours post-dose after 2 weeks of treatment. The clinical relevance of this finding has not been established. The mean ratio of CSF/serum lamivudine concentrations 2 to 4 h after oral administration was approximately 12%. The true extent of CNS penetration of lamivudine and its relationship with any clinical efficacy is unknown.

Metabolism: Dolutegravir is primarily metabolized via UGT1A1 with a minor CYP3A component (9.7% of total dose administered in a human mass balance study). Dolutegravir is the predominant circulating compound in plasma; renal elimination of unchanged drug is low (<1% of the dose). Metabolism of lamivudine is a minor route of elimination. Lamivudine is predominately cleared unchanged by renal excretion. The likelihood of metabolic interactions with lamivudine is low due to the small extent of hepatic metabolism (less than 10%).

Elimination: Dolutegravir has a terminal half-life of approximately 14 hours and an apparent clearance (CL/F) of 0.9-1.05 L/hr based on population pharmacokinetic analyses.

Fifty-three percent of total oral dose is excreted unchanged in the faeces. It is unknown if all or part of this is due to unabsorbed drug or biliary excretion of the glucuronidate conjugate, which can be further degraded to form the parent compound in the gut lumen.

The observed lamivudine half life of elimination is 18 to 19 hours. For patients receiving lamivudine 300 mg once daily, the terminal intracellular half-life of lamivudine-TP was 16 to 19 hours. The mean systemic clearance of lamivudine is approximately 0.32 L/h/kg, predominantly by renal clearance (> 70%) via the organic cationic transport system.

Special Populations and Conditions

Pediatrics: DOVATO has not been studied in the pediatric population.

Dolutegravir: In a pediatric study including 23 antiretroviral treatment-experienced HIV-1 infected adolescents aged 12 to 18 years of age, the pharmacokinetics of dolutegravir was evaluated in 10 adolescents and showed that dolutegravir 50 mg once daily dosage resulted in dolutegravir exposure in pediatric subjects comparable to that observed in adults who received dolutegravir 50 mg once daily ([Table 10](#)).

Table 10 Pediatric pharmacokinetic parameters (n=10)

Age/weight	Dolutegravir Dose	Dolutegravir Pharmacokinetic Parameter Estimates Geometric Mean (CV%)		
		AUC ₍₀₋₂₄₎ μg.hr/mL	C _{max} μg/mL	C ₂₄ μg/mL
12 to <18 years ≥ 40 kg ^a	50 mg once daily ^a	46 (43)	3.49 (38)	0.90 (59)

^a One subject weighing 37 kg received 35 mg once daily.

Lamivudine: Limited data are available in adolescents receiving a daily dose of 300 mg of lamivudine. Pharmacokinetic parameters are comparable to those reported in adults.

Geriatrics: Population pharmacokinetic analysis of dolutegravir using data in HIV-1 infected adults showed that there was no clinically relevant effect of age on dolutegravir exposure. Pharmacokinetic data for dolutegravir and lamivudine in subjects of >65 years old are limited.

Sex: Population pharmacokinetic analyses revealed no clinically relevant effect of gender on the exposure of dolutegravir. No clinically relevant differences in the pharmacokinetics of lamivudine have been observed between men and women.

Pregnancy and Breast-feeding:

The pharmacokinetics of lamivudine during late pregnancy were similar to that of non-pregnant adults. In humans, consistent with passive transmission of lamivudine across the placenta, lamivudine concentrations in infant serum at birth were similar to those in maternal and cord serum at delivery.

There are limited pharmacokinetic data on the use of dolutegravir in pregnancy.

Genetic Polymorphism:

In a meta-analysis using pharmacogenomics samples collected in clinical studies in healthy subjects, subjects with UGT1A1 (n=7) genotypes conferring poor dolutegravir metabolism had a 32% lower clearance of dolutegravir and 46% higher AUC compared with subjects with genotypes associated with normal metabolism via UGT1A1 (n=41).

Ethnic origin:

Population PK analyses using pooled pharmacokinetic data from adult studies revealed no clinically relevant effect of race on the exposure of dolutegravir.

Hepatic Insufficiency:

Pharmacokinetic data has been obtained for dolutegravir and lamivudine alone.

Data obtained for lamivudine in patients with moderate to severe hepatic impairment and for dolutegravir in patients with moderate hepatic impairment show that the pharmacokinetics are not significantly affected by hepatic dysfunction. Dolutegravir is primarily metabolized and

eliminated by the liver. In a study comparing 8 subjects with moderate hepatic impairment (Child-Pugh score B) to 8 matched healthy adult controls, exposure of dolutegravir from a single 50 mg dose was similar between the two groups. The effect of severe hepatic impairment (Child-Pugh score C) on the pharmacokinetics of dolutegravir has not been studied.

Renal Insufficiency:

Pharmacokinetic data have been obtained for dolutegravir and lamivudine alone. DOVATO should not be used in patients with creatinine clearance of less than 30 mL/min because, whilst no dosage adjustment of dolutegravir is necessary in patients with renal impairment, dose reduction is required for the lamivudine component.

Studies with lamivudine show that plasma concentrations (AUC) are increased in patients with renal dysfunction due to decreased clearance.

Renal clearance of unchanged drug is a minor pathway of elimination for dolutegravir. A study of the pharmacokinetics of dolutegravir was performed in subjects (n = 8) with severe renal impairment (CL_{cr} <30 mL/min). No clinically important pharmacokinetic differences between subjects with severe renal impairment (CL_{cr} <30 mL/min) and matching healthy subjects were observed. Dolutegravir AUC, C_{max}, and C₂₄ were lower by 40%, 23%, and 43%, respectively, in subjects with severe renal impairment as compared with matched healthy controls.

Hepatitis B or Hepatitis C Co-infection:

Population pharmacokinetic analysis indicated that hepatitis C virus co-infection had no clinically relevant effect on the exposure to dolutegravir. There are limited pharmacokinetic data on hepatitis B co-infection.

11 STORAGE, STABILITY AND DISPOSAL

Store DOVATO up to 30°C.

Healthcare professionals should recommend that their patients return all unused medications to a pharmacy for proper disposal.

12 SPECIAL HANDLING INSTRUCTIONS

There are no special requirements for use or handling of this product.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Dolutegravir


Drug Substance

Common name: dolutegravir sodium

Chemical name: sodium (4*R*,12*aS*)-9-[[2,4-difluorophenyl)methyl]carbamoyl]-4-methyl-6,8-dioxo-3,4,6,8,12,12*a*-hexahydro-2*H*-pyrido[1',2':4,5]pyrazino[2,1-*b*][1,3]oxazin-7-olate

Molecular formula and molecular mass: C₂₀H₁₈F₂N₃NaO₅
441.36 g/mol

Structural formula:



Physicochemical properties: Dolutegravir sodium is a white to light yellow powder and is slightly soluble in water.

Lamivudine


Drug Substance

Proper name: lamivudine

Chemical name: 2(1*H*)-Pyrimidinone, 4-amino-1-[2-(hydroxymethyl)-1,3-oxathiolan-5-yl]-(2*R*-cis)-

Molecular formula and molecular mass: C₈H₁₁N₃O₃S
229.3

Structural formula:



Physicochemical properties: Lamivudine is a white to off-white crystalline solid with a melting point of 176°C and is soluble in water.

14 CLINICAL TRIALS

14.1 Trial Design and Study Demographics

The efficacy of DOVATO is supported by data from two randomized, double-blind, controlled trials (GEMINI-1 [204861] and GEMINI-2 [205543]) in HIV-1–infected treatment naïve adults, and data from a randomized, open-label, controlled trial (TANGO [204862]) in virologically suppressed HIV-1–infected adults.

Treatment-Naïve Patients

GEMINI-1 and GEMINI-2 are identical 148-week, Phase III, randomized, double-blind, multicenter, parallel-group, non-inferiority controlled trials. A total of 1433 HIV-1 infected antiretroviral treatment-naïve adult subjects received treatment in the trials. Subjects were enrolled with a screening plasma HIV-1 RNA of 1000 c/mL to \leq 500,000 c/mL. Subjects were randomized to a two-drug regimen of TIVICAY (dolutegravir 50 mg) plus 3TC (lamivudine 300mg) administered once daily or to a three-drug regimen TIVICAY (dolutegravir 50 mg) plus TRUVADA (tenofovir/emtricitabine 200mg/300mg) administered once daily. The primary efficacy endpoint for each GEMINI trial was the proportion of subjects with plasma HIV-1 RNA $<$ 50 copies/mL at Week 48 (Snapshot algorithm for the ITT-E population).

The demographic baseline characteristics were similarly distributed between treatment arms (see [Table 11](#)).

Table 11 Summary of Baseline Characteristics for Studies GEMINI-1, GEMINI-2, and Pooled Data (ITT-E Population)

	GEMINI-1		GEMINI-2		POOLED	
	TIVICAY + 3TC	TIVICAY + TRUVADA	TIVICAY + 3TC	TIVICAY + TRUVADA	TIVICAY + 3TC	TIVICAY + TRUVADA
	N=356 (%)	N=358 (%)	N=360 (%)	N=359 (%)	N=716 (%)	N=717 (%)
Baseline HIV-1 RNA (c/mL)						
<100,000	282 (79)	282 (78)	294 (82)	282 (79)	576 (80)	564 (79)
≥100,000 to <500,000	70 (20)	69 (19)	57(16)	69 (19)	127 (18)	138 (19)
≥500,000	4 (1)	7 (2)	9 (3)	8 (2)	13 (2)	15 (2)
Baseline CD4+ (log₁₀ cells/mm³)						
Median	427.0	435.5	427.5	442.0	427.0	438.0
Min., Max.	19, 1399	19, 1305	19, 1364	19, 1497	19, 1399	19, 1497
Baseline CD4+ (cells/mm³), n (%)						
<50	5 (1)	4 (1)	3 (<1)	5 (1)	8 (1)	9 (1)
50 to <200	26 (7)	25 (7)	29 (8)	20 (6)	55 (8)	45 (6)
200 to <350	92 (26)	79 (22)	87 (24)	87 (24)	179 (25)	166 (23)
350 to <500	99 (28)	120 (34)	105 (29)	108 (30)	204 (28)	228 (32)
≥500	134 (38)	130 (36)	136 (38)	139 (39)	270 (38)	269 (38)
Age (y) median (range)	32.0 (18-69)	33.0 (18-66)	32.0 (18-72)	33.0 (18-70)	32.0 (18-72)	33.0 (18-70)
Sex						
Female	59 (17)	52 (15)	54 (15)	46 (13)	113 (16)	98 (14)
Male	297 (83)	306 (85)	306 (85)	313 (87)	603 (84)	619 (86)
Race, n (%)						
American Indian or Alaska Native	31 (9)	28 (8)	21 (6)	29 (8)	52 (7)	57 (8)
Asian	37 (10)	42 (12)	34 (9)	30 (8)	71 (10)	72 (10)
Black/African American	39 (11)	36 (10)	51 (14)	35 (10)	90 (13)	71 (10)
Native Hawaiian or other Pacific Islander	2 (<1)	0	0	1 (<1)	2 (<1)	1 (<1)
White	244 (69)	247 (69)	240 (67)	252 (70)	484 (68)	499 (70)
Multiple Heritage	3 (<1)	5 (1)	14 (4)	12 (3)	17 (2)	17 (2)
Hepatitis B & C Test Results						
B only	0	0	0	0	0	0
C only	26 (7)	28 (8)	13 (4)	21 (6)	39 (5)	49 (7)
B and C	0	0	0	0	0	0
Neither	329 (92)	330 (92)	347 (96)	338 (94)	676 (94)	668 (93)
Missing	1 (<1)	0	0	0	1 (<1)	0
CDC Category						
Stage 0	1 (<1)	0	0	1 (<1)	1 (<1)	1 (<1)
Stage 1	128 (36)	126 (35)	129 (36)	137 (38)	257 (36)	263 (37)
Stage 2	194 (54)	204 (57)	198 (55)	189 (53)	392 (55)	393 (55)
Stage 3	33 (9)	28 (8)	33 (9)	32 (9)	66 (9)	60 (8)

TIVICAY = dolutegravir, 3TC = lamivudine, TRUVADA = tenofovir disoproxil fumarate/emtricitabine

Virologically Suppressed Patients

The efficacy of DOVATO in HIV-infected, antiretroviral therapy experienced, virologically suppressed subjects is supported by data from a Phase III, randomized, open-label, multicenter, parallel-group, non-inferiority controlled trial (TANGO). A total of 741 adult HIV-1 infected subjects who were on a stable suppressive tenofovir alafenamide based regimen (TBR) received treatment. Subjects were randomized in a 1:1 ratio to receive DOVATO once daily or continue with TBR. Randomization was stratified by baseline third agent class (protease inhibitor [PI], integrase inhibitor [INSTI], or non-nucleoside reverse transcriptase inhibitor [NNRTI]). The primary efficacy endpoint was the proportion of subjects with plasma HIV-1 RNA ≥ 50 c/mL (virologic non-response) as per the FDA Snapshot category at Week 48 (Snapshot algorithm adjusting for randomization stratification factor: Baseline Third Agent Class [INSTI, PI, NNRTI]).

At baseline the median age of subjects was 39 years, 8% were female and 21% non-white, 5% were CDC Stage 3 (AIDS) and 98% subjects had Baseline CD4+ cell count ≥ 200 cells/mm³; these characteristics were similar between treatment arms. Subjects had been on ART for a median of 2.8 years and 2.9 years prior to Day 1 for the DOVATO and TBR arms, respectively. Most subjects were on INSTI-based TBR, 78% and 80% in the DOVATO and TBR arms, respectively.

14.2 Study Results

Treatment-Naïve Patients

TIVICAY plus 3TC remained non-inferior to TIVICAY plus TRUVADA through 96 weeks in GEMINI-1 and GEMINI-2 studies. This was supported by the pooled analysis, see [Table 12](#).

Table 12 Virologic Outcomes of Randomized Treatment at Week 48 and Week 96 in GEMINI studies (Snapshot Algorithm, ITT-E Population)

	GEMINI-1				GEMINI-2				GEMINI-1 and GEMINI-2 Pooled Data*			
	TIVICAY + 3TC (N=356) n (%)		TIVICAY + TRUVADA (N=358) n (%)		TIVICAY + 3TC (N=360) n (%)		TIVICAY + TRUVADA (N=359) n (%)		TIVICAY + 3TC (N=716) n (%)		TIVICAY + TRUVADA (N=717) n (%)	
	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96
HIV-1 RNA <50 copies/mL	(320/356) 90%	(300/356) 84%	(332/358) 93%	(320/358) 89%	(335/360) 93%	(316/360) 88%	(337/359) 94%	(322/359) 90%	(655/716) 91%	(616/716) 86%	(669/717) 93%	(642/717) 90%
Treatment Difference[†] (95% confidence intervals)	Week 48: -2.6% (95% CI: -6.7%, 1.5%) Week 96: -4.9% (95% CI: -9.8%, 0.0%)				Week 48: -0.7% (95% CI: -4.3%, 2.9%) Week 96: -1.8% (95% CI: -6.4%, 2.7%)				Week 48: -1.7% (95% CI: -4.4%, 1.1%) Week 96: -3.4% (95% CI: -6.7%, 0.0%)			
Virologic non response	13 (4%)	11 (3%)	6 (2%)	5 (1%)	7 (2%)	11 (3%)	7 (2%)	9 (3%)	20 (3%)	22 (3%)	13 (2%)	14 (2%)
Reasons												
Data in window and ≥50 copies/mL	6 (2%)	1 (<1%)	3 (<1%)	0	2 (<1%)	3 (<1%)	2 (<1%)	4 (1%)	8 (1%)	4 (<1%)	5 (<1%)	4 (<1%)
Discontinued for lack of efficacy	3 (<1%)	4 (1%)	1 (<1%)	2 (<1%)	2 (<1%)	5 (1%)	1 (<1%)	1 (<1%)	5 (<1%)	9 (1%)	2 (<1%)	3 (<1%)
Discontinued for other reasons and ≥50 copies/mL	3 (<1%)	5 (1%)	2 (<1%)	3 (<1%)	2 (<1%)	2 (<1%)	3 (<1%)	3 (<1%)	5 (<1%)	7 (<1%)	5 (<1%)	6 (<1%)
Change in ART	1 (<1%)	1 (<1%)	0	0	1 (<1%)	1 (<1%)	1 (<1%)	1 (<1%)	2 (<1%)	2 (<1%)	1 (<1%)	1 (<1%)
No virologic data at Week 48 window	23 (6%)	45 (13%)	20 (6%)	33 (9%)	18 (5%)	33 (9%)	15 (4%)	28 (8%)	41 (6%)	78 (11%)	35 (5%)	61 (9%)
Reasons												
Discontinued study due to adverse event or death	5 (1%)	12 (3%)	9 (3%)	13 (4%)	5 (1%)	10 (3%)	4 (1%)	8 (2%)	10 (1%)	22 (3%)	13 (2%)	21 (3%)
Discontinued study for other reasons	17 (5%)	33 (9%)	11 (3%)	20 (6%)	12 (3%)	23 (6%)	11 (3%)	18 (5%)	29 (4%)	56 (8%)	22 (3%)	38 (5%)
Missing data during window but on study	1 (<1%)	0	0	0	1 (<1%)	0	0	2 (<1%)	2 (<1%)	0	0	2 (<1%)

	GEMINI-1				GEMINI-2				GEMINI-1 and GEMINI-2 Pooled Data*			
	TIVICAY + 3TC (N=356) n (%)		TIVICAY + TRUVADA (N=358) n (%)		TIVICAY + 3TC (N=360) n (%)		TIVICAY + TRUVADA (N=359) n (%)		TIVICAY + 3TC (N=716) n (%)		TIVICAY + TRUVADA (N=717) n (%)	
	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96
HIV-1 RNA <50 copies/mL by baseline covariates												
									n/N (%)		n/N (%)	
Baseline Plasma Viral Load (copies/mL)												
≤100,000	255 / 282 (90%)	240 / 282 (85%)	263 / 282 (93%)	253 / 282 (90%)	271 / 294 (92%)	259 / 294 (88%)	268 / 282 (95%)	257 / 282 (91%)	526 / 576 (91%)	499 / 576 (87%)	531 / 564 (94%)	510 / 564 (90%)
>100,000	65 / 74 (88%)	60 / 74 (81%)	69 / 76 (91%)	67 / 76 (88%)	64 / 66 (97%)	57 / 66 (86%)	69 / 77 (90%)	65 / 77 (84%)	129 / 140 (92%)	117 / 140 (84%)	138 / 153 (90%)	132 / 153 (86%)
Baseline CD4+ (cells/ mm³)												
≤200	25 / 31 (81%)	20 / 31 (65%)	26 / 29 (90%)	26 / 29 (90%)	25 / 32 (78%)	23 / 32 (72%)	25 / 26 (96%)	22 / 26 (85%)	50 / 63 (79%) ^a	43 / 63 (68%) ^b	51 / 55 (93%)	48 / 55 (87%)
>200	295 / 325 (91%)	280 / 325 (86%)	306 / 329 (93%)	294 / 329 (89%)	310 / 328 (95%)	293 / 328 (89%)	312 / 333 (94%)	300 / 333 (90%)	605 / 653 (93%)	573 / 653 (88%)	618 / 662 (93%)	594 / 662 (90%)
Gender												
Male	268 / 297 (90%)	251 / 297 (85%)	283 / 306 (92%)	274 / 306 (90%)	287 / 306 (94%)	272 / 306 (89%)	297 / 313 (95%)	283 / 313 (90%)	555 / 603 (92%)	523 / 603 (87%)	580 / 619 (94%)	557 / 619 (90%)
Female	52 / 59 (88%)	49 / 59 (83%)	49 / 52 (94%)	46 / 52 (88%)	48 / 54 (89%)	44 / 54 (81%)	40 / 46 (87%)	39 / 46 (85%)	100 / 113 (88%)	93 / 113 (82%)	89 / 98 (91%)	85 / 98 (87%)
Race												
White	220 / 244 (90%)	209 / 244 (86%)	231 / 247 (94%)	223 / 247 (90%)	231 / 240 (96%)	220 / 240 (92%)	242 / 252 (96%)	230 / 252 (91%)	451 / 484 (93%)	429 / 484 (89%)	473 / 499 (95%)	453 / 499 (91%)
African-American/African Heritage/Asian/Other	100 / 112 (89%)	91 / 112 (81%)	101 / 111 (91%)	97 / 111 (87%)	104 / 120 (87%)	96 / 120 (80%)	95 / 107 (89%)	92 / 107 (86%)	204 / 232 (88%)	187 / 232 (81%)	196 / 218 (90%)	189 / 218 (87%)
Age (years)												
<50	294 / 327 (90%)	275 / 327 (84%)	292 / 312 (94%)	280 / 312 (90%)	303 / 324 (94%)	286 / 324 (88%)	305 / 325 (94%)	292 / 325 (90%)	597 / 651 (92%)	561 / 651 (86%)	597 / 637 (94%)	572 / 637 (90%)
≥50	26 / 29 (90%)	25 / 29 (86%)	40 / 46 (87%)	40 / 46 (87%)	32 / 36 (89%)	30 / 36 (83%)	32 / 34 (94%)	30 / 34 (88%)	58 / 65 (89%)	55 / 65 (85%)	72 / 80 (90%)	70 / 80 (88%)

	GEMINI-1				GEMINI-2				GEMINI-1 and GEMINI-2 Pooled Data*			
	TIVICAY + 3TC (N=356) n (%)		TIVICAY + TRUVADA (N=358) n (%)		TIVICAY + 3TC (N=360) n (%)		TIVICAY + TRUVADA (N=359) n (%)		TIVICAY + 3TC (N=716) n (%)		TIVICAY + TRUVADA (N=717) n (%)	
	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96	Week 48	Week 96
<p>TIVICAY = dolutegravir, 3TC = lamivudine, TRUVADA = tenofovir disoproxil fumarate/emtricitabine</p> <p>* The results of the pooled analysis are in line with those of the individual studies, for which the primary endpoint of non-inferiority (in proportion <50 copies/mL plasma HIV-1 RNA at Week 48 based on the Snapshot algorithm for TIVICAY plus 3TC versus TIVICAY plus TRUVADA) was met. The adjusted difference was -2.6 (95% CI: -6.7; 1.5) for GEMINI-1 and -0.7 (95% CI: -4.3; 2.9) for GEMINI-2 with a prespecified non-inferiority margin of -10%.</p> <p>† Based on CMH-stratified analysis adjusting for the following baseline stratification factors: Plasma HIV-1 RNA ($\leq 100,000$ c/mL vs. $> 100,000$ c/mL) and CD4+ cell count (≤ 200 cells/mm³ vs. > 200 cells/mm³). Pooled analysis also stratified by study. Assessed using a non-inferiority margin of -10%.</p> <p>^a With the exception of one patient treated with TIVICAY plus 3TC in GEMINI-1 who was withdrawn due to confirmed loss of virologic response, none of the patients treated with TIVICAY plus 3TC who did not have HIV-1 RNA < 50 copies/mL at Week 48 (based on Snapshot Algorithm) were discontinued for treatment-related reasons by Week 48. A patient in GEMINI-1 whose last HIV-1 RNA was 64,366 copies per/mL was lost to follow up.</p> <p>^b Four subjects treated with TIVICAY plus 3TC were withdrawn for treatment-related reasons (3 due to confirmed loss of virologic response and 1 due to drug-related adverse reactions). Two subjects also had HIV-1 RNA ≥ 50 copies/mL at Week 96 but remained in the study. The other 14 subjects treated with TIVICAY plus 3TC who did not have HIV-1 RNA <50 copies/mL at Week 96 (based on Snapshot algorithm) were discontinued for non-treatment-related reasons by Week 96.</p> <p>N = Number of subjects in each treatment group</p>												

The adjusted mean change from baseline in CD4+ cell count based on the pooled analysis at Week 48 was 224 cells/mm³ for the group receiving TIVICAY + 3TC, and 217 cells/mm³ for the TIVICAY + TRUVADA group. The adjusted mean change from baseline in CD4+ cell count based on the pooled analysis at Week 96 was 269 cells/mm³ for the group receiving TIVICAY + 3TC, and 259 cells/mm³ for the TIVICAY + TRUVADA group.

Virologic outcomes by baseline CD4+ (cells/mm³) in GEMINI-1 and GEMINI-2 are shown in [Table 12](#). In both studies, lower response rates (HIV-1 RNA <50 copies/mL) were observed in patients with baseline CD4+ ≤ 200 cells/mm³. These findings were seen irrespective of baseline plasma HIV-1 RNA.

Virologically Suppressed Patients

The primary analysis in TANGO demonstrated that DOVATO is non-inferior to TBR, with <1% of subjects in both arms experiencing virologic failure (HIV-1 RNA ≥50 c/mL) at Week 48 based on the Snapshot algorithm ([Table 13](#)).

Table 13 Virologic Outcomes of Randomized Treatment of TANGO at Week 48 (Snapshot algorithm)

	DTG/3TC FDC (N=369) n (%)	TBR (N=372) n (%)
Virologic non response (≥ 50 copies/mL)** Treatment Difference[†] (95% confidence intervals)	1 (<1%)	2 (<1%)
Reasons		
Data in window and ≥ 50 copies/mL	0	0
Discontinued for lack of efficacy	0	2 (<1%)
Discontinued for other reasons and ≥ 50 copies/mL	1 (<1%)	0
Change in ART	0	0
HIV-1 RNA <50 copies/mL	(344/369) 93%	(346/372) 93%
No virologic data at Week 48 window	24 (6%)	24 (6%)
Reasons		
Discontinued study due to adverse event or death	12 (3%)	1 (<1%)
Discontinued study for other reasons	12 (3%)	22 (6%)
Missing data during window but on study	0	1 (<1%)
HIV-1 RNA <50 copies/mL by baseline covariates	n/N (%)	n/N (%)
Baseline CD4+ (cells/ mm³)		
<500	92 / 98 (94%)	68 / 74 (92%)
≥ 500	252 / 271 (93%)	278 / 298 (93%)
Baseline Third Agent Class		
NNRTI	49 / 51 (96%)	42 / 48 (88%)
INSTI	268 / 289 (93%)	276 / 296 (93%)
PI	27 / 29 (93%)	28 / 28 (100%)
Gender		
Male	323 / 344 (94%)	319 / 339 (94%)
Female	21 / 25 (84%)	27 / 33 (82%)
Race		
White	279 / 297 (94%)	272 / 289 (94%)
African-American/African Heritage/Other	65 / 72 (90%)	74 / 83 (89%)
Age (years)		
<50	271 / 290 (93%)	260 / 280 (93%)
≥ 50	73 / 79 (92%)	86 / 92 (93%)
TBR = tenofovir alafenamide based regimen; INSTI = Integrase inhibitor; NNRTI = Non-nucleoside reverse transcriptase inhibitor; PI = Protease Inhibitor [†] Based on CMH-stratified analysis adjusting for Baseline third agent class (PI, NNRTI, INSTI). **Based on a 4% non-inferiority margin, DOVATO is non-inferior to TBR at Week 48 in the primary analysis (proportion of subjects with plasma HIV-1 RNA ≥ 50 c/mL) because the upper bound of the 95% CI for the adjusted treatment difference is less than 4% N = Number of subjects in each treatment group		

In TANGO, treatment outcomes between treatment arms were similar across the stratification factor, baseline third-agent class (PI, INSTI, or NNRTI), and across subgroups by age, sex, race, baseline CD4+ cell count, CDC HIV disease stage, and countries. The median change from baseline in CD4+ count at Week 48 was 22.5 cells/mm³ in subjects who received DOVATO and 11.0 cells/mm³ in subjects who received the TBR.

Adolescents

There are no clinical study data with DOVATO in the adolescent population. However, the safety and efficacy of DOVATO in adolescents 12 years of age and older, and weighting at least 40 kg, is supported by the clinical trial data from prior adolescent studies of dolutegravir or lamivudine available in the respective TIVICAY, 3TC and TRIUMEQ product monographs, and also by clinical data from the GEMINI trials with dolutegravir plus lamivudine in adults.

14.3 Comparative Bioavailability Studies

A single-dose, randomized, open-label, 2-period, 2-sequence crossover study was conducted in healthy, adult male and female volunteers (n=76; 50 males and 26 females) to evaluate the comparative bioavailability of an oral 1 x DOVATO (50 mg dolutegravir/300 mg lamivudine) fixed dose combination tablet versus concurrent oral administration of 1 x Dolutegravir 50 mg tablet and EPIVIR (lamivudine 300 mg) tablet under fasting conditions. The effect of a high-fat, high-calorie meal on the bioavailability of the fixed dose combination tablet was also evaluated in a sub-set of the volunteers (n=16; 10 males and 6 females). The comparative bioavailability results from 74 completed subjects (49 males and 25 females) are summarized in tabular format below.

Table 14 Summary of the Comparative Bioavailability Data for Dolutegravir

Dolutegravir (1 x 50 mg) FASTED CONDITIONS From measured data				
Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test¹	Reference²	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (µg.h/mL)	52.3 54.8 (30.5)	45.2 48.4 (36.9)	115.8	(107.2, 125.1)
AUC _I (µg.h/mL)	54.6 57.2 (31.2)	47.2 50.8 (37.8)	115.5	(107.0, 124.7)
C _{max} (µg/mL)	2.91 3.04 (28.7)	2.55 2.70 (32.1)	114.1	(105.3, 123.6)
T _{max} ³ (h)	2.50 (0.500, 6.00)	2.50 (0.500, 5.01)		
T _½ (h) ⁴	15.2 (18.1)	15.4 (17.8)		

- 1 DOVATO (50 mg dolutegravir/300 mg lamivudine) fixed-dose combination tablets.
- 2 Dolutegravir 50 mg tablet and lamivudine 300 mg tablet, administered concurrently.
- 3 Expressed as median (range) only.
- 4 Expressed as the arithmetic mean (CV%) only.

Table 15 Summary of the Comparative Bioavailability Data for Lamivudine

Lamivudine (1 x 300 mg) FASTED CONDITIONS From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	Test ¹	Reference ²	% Ratio of Geometric Means	90% Confidence Interval
AUC _T (µg.h/mL)	13.4 13.6 (17.9)	12.5 12.7 (18.9)	107.0	(104.6, 109.5)
AUC _I (µg.h/mL)	13.6 ³ 13.8 (17.6)	12.8 ³ 13.0 (18.2)	106.4	(104.2, 108.7) ³
C _{max} (µg/mL)	3.22 3.44 (28.4)	2.44 2.53 (26.7)	131.8	(126.2, 137.6)
T _{max} ⁴ (h)	1.00 (0.500, 3.50)	1.00 (0.500, 4.00)		
T _½ (h) ⁵	19.5 (31.1)	20.1 (33.5)		

1 DOVATO (50 mg dolutegravir/300 mg lamivudine) fixed-dose combination tablets.

2 Dolutegravir 50 mg tablet and lamivudine 300 mg tablet, administered concurrently.

3 n=73. One subject was excluded from the statistical analysis of AUC_I because >20% of AUC_I was extrapolated and λz time duration <2x calculated t_{1/2}.

4 Expressed as median (range).

5 Expressed as the arithmetic mean (CV%) only.

15 MICROBIOLOGY

Antiviral Activity in cell culture

Dolutegravir

Dolutegravir exhibited antiviral activity against laboratory strains of wild-type HIV-1 with mean EC₅₀ values of 0.51 nM to 2.1 nM in peripheral blood mononuclear cells (PBMCs) and MT-4 cells.

In a viral integrase susceptibility assay using the integrase coding region from 13 clinically diverse clade B isolates, dolutegravir demonstrated antiviral potency similar to reference laboratory strains, with a mean EC₅₀ of 0.52 nM. When tested in PBMC assays against a panel consisting of 24 HIV-1 clinical isolates [group M (clade A, B, C, D, E, F and G) and group O] and 3 HIV-2 clinical isolates, the geometric mean EC₅₀ was 0.20 nM and EC₅₀ values ranged from 0.02 to 2.14 nM for HIV-1, while the geometric mean EC₅₀ was 0.18 nM and EC₅₀ values ranged from 0.09 to 0.61 nM for HIV-2 isolates.

Lamivudine

The antiviral activity of lamivudine against HIV-1 was assessed in a number of cell lines including monocytes and PBMCs using standard susceptibility assays. EC₅₀ values were in the range of 0.003 µM to 2 µM (1 µM = 0.23 µg/mL). The EC₅₀ values of lamivudine against different HIV-1 clades (A to G) ranged from 0.001 to 0.120 µM, and against HIV-2 isolates from 0.002 to 0.041 µM in PBMCs. Ribavirin (50µM) decreased the anti-HIV-1 activity of lamivudine by 3.5-fold in MT-4 cells.

Antiviral Activity in combination with other antiviral agents

Dolutegravir

The following drugs were not antagonistic with dolutegravir in *in-vitro* assessments conducted in checkerboard format: stavudine, abacavir, efavirenz, nevirapine, lopinavir, amprenavir, enfuvirtide, maraviroc, adefovir and raltegravir. In addition, the anti-HCV drug ribavirin had no apparent effect on dolutegravir activity.

Lamivudine

No antagonistic effects *in vitro* were seen with lamivudine and other antiretrovirals (tested agents: abacavir, didanosine, nevirapine, zalcitabine, and zidovudine).

Effect of Human Serum and Serum Proteins

In vitro studies suggested a 75-fold shift in EC₅₀ of dolutegravir in the presence of 100% human serum (by method of extrapolation), and the protein adjusted EC₉₀ (PA-EC₉₀) in PBMCs for dolutegravir was estimated to be 0.064 µg/mL. Dolutegravir trough concentration for a single 50 mg dose in integrase inhibitor naïve patients was 1.20 µg/mL, 19 times higher than the estimated PA-EC₉₀. Lamivudine exhibits linear pharmacokinetics over the therapeutic dose range and displays low plasma protein binding (less than 36%).

Resistance in vitro and in vivo (dolutegravir)

Isolation from wild type HIV-1 and activity against resistant strains

Viruses highly resistant to dolutegravir were not observed during the 112 day passage of strain III B, with a 4.1-fold maximum fold change (FC) observed for the passaged resistant virus populations with substitutions at the conserved IN positions S153Y and S153F.

Passage of the wild type HIV-1 strain NL432 in the presence of dolutegravir selected for E92Q (passage population virus FC=3.1) and G193E (passage population virus FC=3.2) substitutions on Day 56. Additional passage of wildtype subtype B, C, and A/G viruses in the presence of dolutegravir selected for R263K (site-directed mutant FC = 1.5), G118R (site-directed mutant FC = 10), and S153T.

Treatment-naïve HIV-1 infected subjects receiving dolutegravir

No INI-resistant mutations or treatment emergent resistance to the NRTI backbone therapy were isolated with dolutegravir 50 mg once daily in treatment-naïve studies.

Resistance in vitro and in vivo (lamivudine)

HIV-1 resistance to lamivudine involves the development of a M184I or M184V amino acid change close to the active site of the viral RT. This variant arises both during *in vitro* selection and in HIV-1 infected patients treated with lamivudine-containing antiretroviral therapy. M184V mutants display greatly reduced susceptibility to lamivudine and show diminished viral replicative capacity *in vitro*.

Resistance in vivo (dolutegravir plus lamivudine)

No subjects that met the protocol-defined confirmed virologic withdrawal (CVW) criteria across the pooled GEMINI-1 and GEMINI-2 studies through Week 96 or in the TANGO study through Week 48 had emergent INSTI or NRTI resistance substitutions.

Cross-resistance

Cross resistance between lamivudine and antiretrovirals from other classes (e.g. protease inhibitors (PI) or non-nucleoside reverse transcriptase inhibitors (NNRTIs)), is unlikely.

Site-directed INSTI mutant virus

Dolutegravir activity was determined against a panel of 60 INSTI-resistant site-directed mutant HIV-1 viruses (28 with single substitutions and 32 with 2 or more substitutions). The single INSTI-resistance substitutions T66K, I151L, and S153Y conferred a greater than 2-fold decrease in dolutegravir susceptibility (range: 2.3-fold to 3.6-fold from reference). Combinations of multiple substitutions T66K/L74M, E92Q/N155H, G140C/Q148R, G140S/Q148H, R or K, Q148R/N155H, T97A/G140S/Q148, and substitutions at E138/G140/Q148 showed a greater than 2-fold decrease in dolutegravir susceptibility (range: 2.5-fold to 21-fold from reference).

Recombinant resistant clinical isolates

Dolutegravir activity was measured for 705 raltegravir resistant recombinant isolates from clinical practice; 93.9% (662/705) of the isolates had a dolutegravir FC ≤10 and 1.8% had a DTG FC >25. Mutants with Y143 and N155 pathway had mean FCs of 1.2 and 1.5, respectively, while Q148 + 1 mutant and Q148 + ≥2 mutants mean FCs were 4.8 and 6.0, respectively.

Cross-resistance conferred by the M184V reverse transcriptase

Cross-resistance is limited within the nucleoside inhibitor class of antiretroviral agents. Zidovudine and stavudine maintain their antiretroviral activities against lamivudine-resistant HIV-1. Abacavir and tenofovir maintain antiretroviral activity against lamivudine-resistant HIV-1 harbouring only the M184V mutation.

16 NON-CLINICAL TOXICOLOGY

Carcinogenesis/mutagenesis

Dolutegravir was not mutagenic or clastogenic using *in vitro* tests in bacteria and cultured mammalian cells, and an *in vivo* rodent micronucleus assay. Dolutegravir was not carcinogenic in long term studies in the mouse and rat.

Lamivudine was not mutagenic in bacterial tests, but like many nucleoside analogues it shows activity in the *in vitro* mammalian tests such as the mouse lymphoma assay. This is consistent with the known activity of other nucleoside analogues. The results from two *in vivo* rat micronucleus tests with lamivudine were negative.

Lamivudine did not show any genotoxic activity in additional *in vivo* studies in rats (metaphase analysis of bone marrow and unscheduled DNA synthesis). The results of long-term carcinogenicity studies in mice and rats did not show any carcinogenic potential at exposures approximately 11 to 65 times higher than human clinical exposure based on AUC.

Reproductive Toxicology

In a peri-/post-natal/juvenile toxicity study with lamivudine in rats, some histological inflammatory changes at the ano-rectal junction and slight diffuse epithelial hyperplasia of the cecum were observed in dams and pups at the high-dose level. An increased incidence of urination upon handling was also seen in some offspring at exposures >50 times the human clinical exposure based on C_{max} . In addition, a reduction in testes weight was observed in juvenile males (at exposures >125 times the human clinical exposure based on C_{max}) which was associated with slight to moderate dilatation of the seminiferous tubules.

Fertility

Fertility studies in the rat have shown that dolutegravir and lamivudine had no effect on male or female fertility. Dolutegravir did not affect male or female fertility in rats at doses up to 1000 mg/kg/day, the highest dose tested (33 times the 50 mg human clinical exposure, based on AUC). Lamivudine did not affect male or female fertility in rats at doses up to 2000 mg/kg BID, the highest dose tested (>90 times the 300 mg human clinical exposure, based on AUC).

17 SUPPORTING PRODUCT MONOGRAPHS

1. 3TC (tablets, 300 mg 150 mg; oral solution, 10 mg/mL; lamivudine), submission control #202946, Product Monograph, ViiV Healthcare ULC. (May 12, 2017)
2. TIVICAY (tablets, 10, 25, and 50 mg dolutegravir), submission control #217790, Product Monograph, ViiV Healthcare ULC. (August 27, 2018).

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE
PATIENT MEDICATION INFORMATION

DOVATO
dolutegravir and lamivudine tablets

Read this carefully before you start taking **DOVATO** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **DOVATO**.

Serious Warnings and Precautions

Worsening of hepatitis B virus in people who have HIV-1 infection

If you have a hepatitis B infection, you should not stop taking **DOVATO** without talking to your doctor. If you have to stop taking **DOVATO** your hepatitis may worsen. Your doctor will monitor your liver function for several months and may give you a new medication to treat your hepatitis B infection.

What is **DOVATO used for?**

- **DOVATO** is used to treat HIV (human immunodeficiency virus) infection in adults and adolescents over the age of 12 years and weighing at least 40 kg.

How does **DOVATO work?**

DOVATO contains two medicines that are used to treat HIV infection: Dolutegravir and Lamivudine.

These medicines work together to reduce the amount of virus in your body and keep it at a low level. This helps maintain the number of CD4+ cell count in your blood. CD4+ cells are a type of white blood cells that are important in helping your body to fight infection. **DOVATO** does not cure HIV infection.

What are the ingredients in **DOVATO?**

Medicinal ingredients: 50 mg dolutegravir (as dolutegravir sodium), 300 mg lamivudine.

Non-medicinal ingredients: hypromellose, macrogol / PEG, magnesium stearate, mannitol (E421), microcrystalline cellulose, povidone (K29/32), sodium starch glycolate, sodium stearyl fumarate, titanium dioxide

****DOVATO** comes in the following dosage forms:**

50 mg dolutegravir / 300 mg lamivudine fixed dose combination tablets.

Do not use **DOVATO if:**

- You are allergic (*hypersensitive*) to dolutegravir (TIVICAY, TRIUMEQ or JULUCA) or lamivudine (3TC, KIVEXA, COMBIVIR, TRIZIVIR or TRIUMEQ) or to any of the other ingredients of **DOVATO**. See "What are the ingredients in **DOVATO**?".
- You are taking dofetilide (to treat heart conditions).
- You are taking fampridine (also known as dalfampridine) used to treat multiple sclerosis.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take DOVATO. Talk about any health conditions or problems you may have, including if you:

- have had kidney or liver problems, including hepatitis B or C infection.
- have ever had a severe skin rash when taking dolutegravir (TIVICAY, TRIUMEQ or JULUCA) or lamivudine (3TC, KIVEXA, COMBIVIR, TRIZIVIR or TRIUMEQ).
- have ever had high levels of acid in the blood (lactic acidosis).
- have ever had an increase in your blood sugar (glucose) or levels of fats (lipids) in your blood.
- you have symptoms of an infection or inflammation, as these may flare up while on HIV treatment or you may have even stronger reactions to new infections than you would normally have.

Other warnings you should know about:

Pregnancy

Talk to your doctor if you are pregnant or plan to become pregnant. Your doctor will consider the benefit to you and the risk to your baby when taking DOVATO while you are pregnant.

- Taking DOVATO at the time of becoming pregnant or during the first 12 weeks of pregnancy may increase the risk of a type of birth defect called neural tube defect such as spina bifida (malformed spinal cord).
- In babies and infants exposed to one of the ingredients in DOVATO during pregnancy or labour, small temporary increases in blood levels of a substance called lactate have been observed. There have also been very rare reports of diseases that affect the nervous system such as delayed development and seizures.
- There is a registry for women who take antiretroviral medicines during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare professional about how you can take part in this registry.
- If you could get pregnant while taking DOVATO, you need to use a reliable method of birth control to prevent pregnancy.

Breastfeeding

Do not breastfeed while taking DOVATO. There is a risk of passing HIV-1 to your baby if you breastfeed. DOVATO can also be passed through breast milk and harm your baby. If you are breastfeeding or planning to breastfeed, talk with your healthcare professional about the best way to feed your baby.

Infecting others with HIV

DOVATO will not stop you from passing HIV to others, although this risk is lower if you take your HIV medicine as instructed by your healthcare professional. You should take steps to avoid this by:

- Using condoms when you have oral or penetrative sex.
- Not reusing or sharing needles, syringes, or other injection equipment.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with DOVATO:

- antacids to treat indigestion and heartburn and laxatives to treat constipation.
 - Some antacids and laxatives can stop DOVATO from being absorbed into your body and not make it work as well.
 - DOVATO should be taken at least 2 hours before or 6 hours after you take an antacid or

- laxative.
- Other acid-lowering medicines like ranitidine and omeprazole can be taken at the same time as DOVATO.
- calcium and iron supplements (non-antacids).
 - Taking these supplements at the same time as DOVATO can stop DOVATO from being absorbed into your body and not make it work as well.
 - DOVATO should be taken at least 2 hours before or 6 hours after you take these supplements.
 - You can take supplements containing calcium or iron at the same time as DOVATO if you take both with food.
- metformin, to treat diabetes
- rifampin, to treat some bacterial infections, such as tuberculosis (TB)
- phenytoin and phenobarbital, to treat epilepsy
- oxcarbazepine and carbamazepine, to treat epilepsy and bipolar disorder
- St. John's wort, (*Hypericum perforatum*), a herbal remedy to treat depression
- sorbitol-containing medicines (usually liquids) used regularly
- trimethoprim/sulfamethoxazole

Talk to your healthcare professional for further advice if you are taking any of these medicines. For some of these medicines, your healthcare professional may need to adjust the dose of one of your medicines in order for it to work properly.

How to take DOVATO:

Always take DOVATO every day exactly as your doctor has told you to. DOVATO can be taken with or without food. Check with your healthcare professional if you're not sure.

Usual dose:

The usual dose of DOVATO in adults and adolescents 12 years of age and older and weighing at least 40 kg is one tablet taken once a day.

Take DOVATO for as long as your doctor recommends. Don't stop unless your doctor advises you to.

Overdose:

If you think you have taken too much DOVATO, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you miss a dose, take DOVATO as soon as you remember. If your next dose is due within 4 hours, skip the dose you missed and take the next one at the usual time. Then continue your treatment as before. Don't take a double dose to make up for a missed dose.

What are possible side effects from using DOVATO?

These are not all the possible side effects you may feel when taking DOVATO. If you experience any side effects not listed here, contact your healthcare professional.

The most common side effects of DOVATO are:

- Headache
- Diarrhea
- Drowsiness
- Feeling sick (nausea)
- Trouble sleeping (insomnia)
- Anxiety

Additional side effects that may occur include: itching (pruritus), any new infection, kidney problems, being sick (vomiting), stomach pain (abdominal pain), stomach (abdominal) discomfort, intestinal gas (flatulence), fever, feeling tired, muscle pain, dizziness, abnormal dreams, feeling sleepy and depression (feelings of deep sadness and unworthiness).

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Allergic (hypersensitivity) reactions: <ul style="list-style-type: none"> • Skin rash, fever, lack of energy, swelling of the mouth or face causing difficulty in breathing, muscle or joint aches 			✓
Blood problems: <ul style="list-style-type: none"> • Low red blood cell count (Anemia) <ul style="list-style-type: none"> ○ lack of energy, breathlessness • Low white blood cell count (neutropenia) <ul style="list-style-type: none"> ○ Increased chance of infection • Low platelets (thrombocytopenia) <ul style="list-style-type: none"> ○ Increased chance of bruising 		✓	
Liver problems and blood test results: <ul style="list-style-type: none"> • Inflammation (Hepatitis) • Bilirubin increase (substance produced by liver) • Increase of muscle enzymes (CPK) • Increase in a kidney function blood test result (creatinine) 		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE			
High level of acid in the blood (Lactic Acidosis) <ul style="list-style-type: none"> • Deep, fast, or hard to catch a breath (hard to breathe) • Numbness/weakness of arms or legs • Swollen/enlarged liver • Feeling sick (nausea), being sick (vomiting), stomach pain 		✓	
Liver failure: <ul style="list-style-type: none"> • Extremely high liver blood test results • Yellowing of the skin and the whites of the eyes • Dark/tea coloured urine • Pale coloured stools/ bowel movements • Nausea/vomiting • Loss of appetite • Pain, aching/tenderness on right side below ribs 		✓	
Suicidal thoughts or actions (mainly in patients who have had depression or mental health problems before)		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

<p>Reporting Side Effects</p> <p>You can report any suspected side effects associated with the use of health products to Health Canada by:</p> <ul style="list-style-type: none"> • Visiting the Web page on Adverse Reaction Reporting (https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html) for information on how to report online, by mail or by fax; or • Calling toll-free at 1-866-234-2345. <p><i>NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.</i></p>
--

Storage:

Store DOVATO up to 30°C.

Keep out of reach and sight of children.

Proper disposal:

Do not throw any medicines away in the garbage, down the sink drain or in the toilet. Give all unused medicines to your local pharmacy for proper disposal. This will help to protect the environment.

If you want more information about DOVATO:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<http://hc-sc.gc.ca/index-eng.php>); the manufacturer's website www.viivhealthcare.ca, or by calling 1-877-393-8448.

This leaflet was prepared by ViiV Healthcare ULC

Last Revised September 15, 2021

*©2021 ViiV Healthcare group of companies or its licensor
Trademarks are owned by or licensed to the ViiV Healthcare group of companies*